

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



ISSN: 09/616,283; Art Unit: 1645  
Attorney Docket No. VRXB-P01-001

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Application of:

GOODNOW

Serial No: 09/616,283

Filed: July 14, 2000

For: SYSTEM FOR DETECTING  
BACTERIA IN BLOOD, BLOOD  
PRODUCTS, AND FLUIDS OF  
TISSUES

Art Unit: 1645

Attorney Docket No. VRXB-P01-001

Examiner: J. Hines

Assistant Commissioner of Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

**Declaration Under 37 C.F.R. §1.132**

Sir:

I, Harvey G. Klein, MD., hereby declare as follows:

1. I am the Chief of the Department of Transfusion Medicine at the Warren C. Magnuson Clinical Center, National Institutes of Health. I have been working and conducting research in the field of transfusion medicine for over thirty (30) years. I serve on the Food and Drug Administration's (FDA) Blood Products Advisory Committee. I actively hold leadership positions in the American Association of Blood Banks, the American Blood Commission, the American Society of Hematology, and the American Society of Apheresis. I currently serve as Chairman of the Committee of Revision for Blood and Blood Products for the U.S. Pharmacopeia. I am a diplomat of the American Board of Internal Medicine and a diplomat of the American Board of Pathology. I have authored and co-

more than 175 publications on transfusion medicine. Accordingly, my curriculum vitae is attached as Appendix A.

2. I have read the above-identified application, the pending claims, and the Office Action mailed on September 29, 2003.
3. I understand that the Examiner has stated that the invention as described and claimed in the above-identified application is obvious in view of the teachings of Fisher et al. WO 98/57994, McLaughlin (U.S. Patent 4,683,196), Tadler et al. (*J. Clin. Lab. Anal.* 3: 21-25 (1989)), Erich et al. (*J. Immunol.* 143(12): 4053-4060, 1989), and Chang et al. (U.S. Patent 5,200,323).
4. For the reasons stated below, I respectfully disagree with the Examiner. I have been working in the field of transfusion medicine for over thirty (30) years and over the past two decades the issue of bacterial contamination has been a source of primary concern for blood collectors and transfusion service scientists and clinicians. It is estimated that as many as one in 12,000 transfusions lead to a severe septic reaction and as many as one in 46,000 transfusions can lead to death due to bacterial contamination. In the recent years, it has been further reported that an unexpectedly large number of deaths were attributed to bacterial contamination/sepsis. For example, the fatality rate for transfusion related bacterial sepsis during the period from 1999 to 2002 averaged more than 15%.<sup>1</sup>

---

(a) <sup>1</sup> People's Choice Award - 2003 FDA Science Forum; "Transfusion Related Fatalities from Bacterial Contamination of Blood Components," L.E.Simmons, MT(ASCP)<sup>1</sup>, M.A.Knippen<sup>2</sup>, L.G.Holness, M.D.<sup>3</sup>, <sup>1</sup>OCBQ, CBER, FDA, Rockville, MD, <sup>2</sup>OCBQ, CBER, FDA, Rockville, MD, <sup>3</sup>OBRR, CBER, FDA, Rockville, MD

Although, blood banks routinely test each unit of donated blood for human immunodeficiency virus (HIV), hepatitis B and C virus, syphilis, and Human T-cell Lymphotropic Virus (HTLV) and discard units which have abnormal test results, they do not test for bacterial contamination because of the lack of a safe and effective test.<sup>2</sup> Accordingly, the nation's leading blood safety experts have been repeatedly calling for immediate action from the blood banking community to initiate a program to detect the presence of bacteria in blood and blood products.

5. As early as 1992, P. Ann Hoppe<sup>3</sup> reported that although numerous studies had been performed by the FDA and others, no rapid and reliable tests existed that could be readily applied in the blood bank setting. See *Transfusion* 1992; 32(3): 199-201.
6. In December 1997, a multi-center study for the systematic collection of data concerning bacterial contamination of blood components (BaCon Study) was initiated under the guidance of the American Association of Blood Banks (AABB), the American Red Cross (ARC), and the Center for Disease Control and prevention (CDC). The goals of this study included determining the rates of bacterial contamination associated with recipient transfusion reactions, identification of responsible micro-organisms, and identification of risk factors for bacterial contamination.
7. On February 23, 1999, Jane Henney, the FDA commissioner testified as follows before the U.S. House of Representatives in a hearing on issues of blood safety: "The safety and

---

<sup>2</sup>American Red Cross web site on safety of donated blood:  
[www.givelife2.org/donor/bloodsafety.asp](http://www.givelife2.org/donor/bloodsafety.asp). See Appendix B

<sup>3</sup> The Acting Director, Division of Transfusion Science of the FDA.




adequacy of the blood supply and blood products is one of the highest priorities of the FDA and the Department of Health and Human Resources. . . For a number of serious and life-threatening infections, there is a limited period after a possible donor has been infected, which the infection is not detectable by available methods. ... The risk to patients from bacterial contamination of blood and from blood bank error must also be reduced.”


8. Despite this long felt need and high level of interest in the field for a rapid and effective screening assay to detect clinically relevant amounts of bacteria in blood and blood products, to the best of my knowledge no such reliable test is currently on the market. In fact, due to the lack of such reliable testing methods, the Food and Drug Administration (FDA) guidelines require shorter storage times as a means for controlling the increase in bacterial contamination upon storage. However, the trade-off is reduced blood availability.
9. I list below certain tests that have been developed over the past two decades in an effort to alleviate this problem. However, none of these have proven to be effective in detecting clinically relevant amounts of bacteria.
  - RNA probes were developed as universal probes to detect bacterial contamination; however, this method was too cumbersome for routine blood bank screening and was associated with a high rate of contamination. These probes are no longer being marketed by the manufacturer.
  - Visual inspections also did not prove to be a useful method for detecting bacterial contamination.

- Testing for glucose levels was not found to be a viable alternative because certain contaminants like *S. epidermis*, a common skin contaminant, grows slowly and glucose changes could not be detected until later in the storage period.
- Filtration methods had no impact on the bacterial levels – in no case were the bacteria eliminated, and in fact, bacteria reached the same titer at the stationary phase of growth, whether or not the pool was filtered.
- Certain antigen detection systems using latex agglutination, fluorescent antibody stains, and enzyme-linked immunosorbent, are available, however, these methods have been plagued with variable sensitivity and specificity. As taught in the Verax application, an obvious concern with regards to detection of blood contaminants is the requirement of a ubiquitous microbial antigen to ensure detection of diverse bacterial species.
- Around the mid-1990s, the American Red Cross, in collaboration with a commercial company Binax, developed a prototype assay using immunological methods to detect bacterial contamination of blood components. The test was developed to detect the presence of a common microbial antigen, peptidoglycan, by immunochromatography. However, Binax subsequently abandoned the test due to its inability to detect all the bacteria implicated in platelet transfusions as peptidoglycans are not expressed on the surface of Gram negative bacteria.

10. Therefore, to date, despite the repeated call for improved testing methodologies, and despite the numerous attempts to develop effective tests, there is no effective method for detecting bacterial contamination. I believe that the Verax test provides a non-obvious solution to meet this long-felt need in the blood banking arena.
11. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title XVIII of the United States Code and that willful false statements may jeopardize the validity of this Application for Patent or any patent issuing thereon.

Dated: 12/1/2003

Signature:   
Dr. Harvey G. Klein  
Chief, Department of Transfusion Medicine  
Warren C. Magnuson Clinical Center  
National Institutes of Health



## **CURRICULUM VITAE**

**HARVEY G. KLEIN, M.D.**

**DATE OF BIRTH:** May 8, 1943

**PLACE OF BIRTH:** Boston, Massachusetts

**MARITAL STATUS:** Married, June 20, 1965

**WIFE'S NAME:** Sigrid Petri

**CHILDREN:** Susanna Rebecca, born July 26, 1969  
Stephan Eliot, born February 3, 1972  
Melissa Kathryn, born June 30, 1976

**HOME ADDRESS:** 13628 Canal Vista Court, Potomac, MD 20854

**OFFICE ADDRESS:** National Institutes of Health  
Warren G. Magnuson Clinical Center  
Department of Transfusion Medicine  
10 Center Drive, Building 10, Room 1C711  
Bethesda, MD 20892-1184

### **EDUCATION:**

|           |  |
|-----------|--|
| 1955-1961 | Boston Latin School (Franklin Medalist)    |
| 1961-1965 | Harvard College, A.B. (magna cum laude)    |
| 1965-1969 | The Johns Hopkins School of Medicine, M.D. |

### **POSTGRADUATE TRAINING:**

|           |   |
|-----------|---|
| 1969-1970 | Internship, The Johns Hopkins Hospital<br>Department of Medicine            |
| 1970-1972 | Resident, The Johns Hopkins Hospital,<br>Department of Medicine             |
| 1972-1973 | Fellow in Hematology, The Johns Hopkins<br>Hospital, Department of Medicine |

### **MILITARY SERVICE:**

|           |   |
|-----------|---|
| 1973-1975 | Commissioned Corps, Public Health Service |
|-----------|---|

## **PROFESSIONAL APPOINTMENTS:**

|              |  |
|--------------|--|
| Present:     | Chief, Department of Transfusion Medicine  |
| 1997-present | Executive Committee, Warren G. Magnuson Clinical Center                                |
| 1990-94      | Special Assistant to the Director for Science, Warren G. Magnuson Clinical Center, NIH |
| 1983         | Tenured Scientist  |
| 1976-83      | Assistant Chief, Blood Bank Department, NIH  |
| 1976-83      | Chief, Blood Services Section, Clinical Center Blood Bank, NIH                         |
| 1975-6       | Acting Chief of Blood Services, Clinical Center Blood Bank, NIH                        |
| 1974-5       | Chief, Hemorrhagic Diseases Program, Division of Blood Diseases, NHLBI, NIH            |
| 1973-5       | Assistant to the Director, Division of Blood Diseases and Resources, NHLBI, NIH        |

## **CERTIFICATION:**

Maryland Medical License, 1969 (#K8711)  
Diplomate, American Board of Internal Medicine, 1972 (#39283)  
Diplomate, Hematology Subspecialty - ABIM, 1976 (#39283)  
Diplomate, American Board of Pathology, Blood Bank Specialty, 1977

## **ACADEMIC APPOINTMENT:**

1995-present Johns Hopkins School of Medicine – Visiting Professor, Pathology  
1990-present Johns Hopkins School of Medicine – Visiting Professor, Medicine

## **PROFESSIONAL SOCIETIES:**

American Federation of Medical Research  
American Society of Hematology  
American Association of Blood Banks  
American Society for Apheresis  
International Society of Blood Transfusion

## **AWARDS AND HONORS:**

Detur Award (Harvard)  
Phi Beta Kappa  
Award of Distinction, American Society of Medical Writers, 1975  
Commendation Medal, Public Health Service, 1980

Peacock Lecturer, University of Texas (Dallas), 1984  
 Equal Opportunity Achievement Award, Public Health Service, 1985  
 Elmer L. DeGowin Lectureship, University of Iowa, 1986  
 Meritorious Service Medal, Public Health Service, 1989  
 PHS Unit Commendation, 1991  
 Clarence Schein Memorial Lectureship, Montefiore Medicine Center,  
 New York, 1991  
 Visiting Professor, Mayo Clinic Section of Transfusion Medicine, 1991  
 NIH Clinical Center Director's Award, 1992  
 John B. Alsever Award (contributions to the blood banking sciences), 1993  
 Annual Lecture Award - American Society for Apheresis and 5th World  
 Apheresis Association Congress, 1994  
 Charles E. Walter Memorial Award, Mid Atlantic Association of Blood Banks,  
 1995  
 Allen Latham Award (European Society for Haemapheresis), 1995  
 Tibor J. Greenwalt Lectureship, 1996  
 Cohn DeLaval Medal (World Apheresis Association), 1996  
 Joseph R. Bove Visiting Professorship, Yale, 1997  
 Rita and Taft Schreiber Memorial Lectureship Cedars-Sinai Medical Center, 1998  
 Corresponding Membership, German Blood Transfusion Society, 1998  
 Distinguished Visiting Professor of Pathology, The Johns Hopkins School of  
 Medicine, 1999  
 Visiting Professor of Pathology, University of Pennsylvania, 1999  
 Chairman, 24<sup>th</sup> International Symposium of Blood Transfusion, The  
 Netherlands, 1999  
 HHS Secretary's Distinguished Services Award, 2000  
 NIH Director's Award, 2000  
 Millennial Medical Festival Lecturer, Royal College of Physicians, London, 2000  
 Distinguished Lecturer, Texas Blood Institute, San Antonio, 2000  
 Emily Cooley Award (American Association of Blood Banks) 2001  
 Francis Morrison Memorial Lecture (American Society for Apheresis) 2003  
 Keynote Speaker, Paul Ehrlich Institute (Frankfurt) 2003

#### **CONSULTANT APPOINTMENT:**

Attending Physician, Clinical Hematology Branch, NHLBI, 1975-96  
 FAES Graduate School at NIH (Immunohematology), 1976-1986  
 National Heart, Lung and Blood Institute, Division of Extramural Affairs,  
 1977-present  
 National Blood Resources Education Program (NBREP) 1988-91  
 Uniformed Services University of Health Science, Clinical Asst. Prof.,  
 1977-present  
 Letterman Army Institute of Research, Blood Peer Review Panel, 1987-92  
 Erythropoietin Advisory Board - Ortho Biotech Corp., 1989-95  
 Haemonetics Corp. Scientific Advisory Board

AIBS - Medical Free-electron Laser Program, 1990-91  
 Somatogen Corp., Advisory Board (recombinant human hemoglobin), 1989-98  
 American Medical Association - Drug Evaluations Annual, 1992  
 Cryopharm - Scientific Advisory Board (lyophilized blood cells), 1994-5  
 Zymequist - Scientific Advisory Board (universal red cells), 1994-present  
 AMA Therapeutic Technology Assessment  
 Office of Naval Research – Navy-Army Blood Program, 1998-2000  
 British Blood Authority, 1998 - present  
 Alliance Pharmaceuticals (perfluorocarbon oxygen carriers), 1998-Present  
 New Jersey Department of Health & Senior Services (Bloodless Surgery) 1999  
 Sangart (hemoglobin-based oxygen carriers), 1999-Present  
 World Health Organization, 2000  
 Viacell Medical Scientific Advisory Board (cell expansion) 2000 – Present  
 Vitex Scientific Advisory Board (pathogen reduction of blood) 2001-Present  
 Gambro BCT Medical Advisory Board (pathogen reduction of blood) 2001  
 Johns Hopkins Hospital Advanced Transfusion Practices Center, 2000 – 2002

#### **EDITORIAL BOARD MEMBERSHIP:**

Editor-in-Chief, Journal of Clinical Apheresis, 1986-94  
 Transfusion, 1994-present  
 Transfusion Medicine Reviews, 1994-present  
 Blood, 1988 - 1993  
 Infusion Therapy and Transfusion Medicine (German Society of Blood Transf.)  
 NIH Alumni Update (Founding Member), 1989 - 1996

#### **PROFESSIONAL ACTIVITIES:**

##### **American Association of Blood Banks**

President, 2000-2001  
 President – elect, 1999-2000  
 Vice President, 1997-1999  
 Board of Directors, 1995-2002  
Chairman, Board of Trustees, Research Foundation, 1985-9  
Co-chairman, Scientific Strategic Plan, 1989  
 Transfusion Practices Committee, 1982-8  
 Therapeutic Hemapheresis Committee, 1983-6  
 Nominations Committee, 1990, 1992, 2003  
 Ad Hoc Standards Committee for Marrow Processing Facilities, 1991  
 Strategic Planning Committee, 1991-3,  
 Standards Committee, 1991-6, Chairman, 1993-6  
 North America Task Force for Stem Cell Standards, 1994-6  
 Board of Directors, National Blood Data Resource Center, 2002-3

## **American Blood Commission**

Board of Directors, 1985-8, ASFA representative

## **American National Red Cross**

Blood Services Advisory Committee, 1981-3

Committee for Protection of Human Subjects, 1983-7

Scientific Council for Biomedical Research , 1985-9

Chairman, Transfusion Science Initial Review Group, 1985-9

Board of Directors, Washington Region , 1989-94

Chairman, Medical Advisory Committee, Blood Services Washington Region, 1988-92

Blood Services Committee, Washington Region, 1988-90

Blood Services Committee, Greater Chesapeake and Potomac Region, 1990

Board of Directors, Chesapeake and Potomac Blood Program, 1992 - present

## **American Society of Hematology**

Scientific Subcommittee in Clinical Laboratory Hematology, 1991-97

Transfusion Medicine Subcommittee, 1993-8

Chairman, Transfusion Medicine Education Program, 1997, 1998

International Outreach Initiative – Subcommittee for Asia

## **American Society for Apheresis**

President, 1985

Board of Directors, 1981-7

Chairman, Clinical Applications Committee, 1983-7

Chairman, Scientific Program, 1984, 1990

## **FDA**

Blood Products Advisory Committee, 2002 - present

Advisory Panel on Orphan Drugs

Blood donors at risk for AIDS - subject matter expert

Workshop on Criteria for Evaluation of Red Cell Substitutes, 1999

Workshop on Leukoreduction of Blood Components, 1999

## **Haemonetics Corporation**

Medical Advisory Board, 1988-99



Board of Directors, 1998-2002

**Institute of Medicine**

Blood Safety Forum, 1994-95  
Committee on Resuscitation Fluids, 1998

**International Society of Blood Transfusion**

Scientific Committee – Int'l Congress (2000) Vienna  
BEST Working Party, 1999-present  
Scientific Committee International Congress (2004 Edinburgh)

**Metropolitan Washington Blood Banks, Inc.**

Board of Directors, 1979-82  
Chairman, Committee on Regionalization, 1979-80

**National Marrow Donor Program (NMDP)**

President, NMDP Council, 1993-4  
Board of Directors, 1992-95  
Membership Committee, 1988-92  
Research and Publications Committee, 1990-95  
International Affairs Committee, 1990-92  
Vice President for Donor Centers, NMDP Council, 1992-94

**National Heart, Lung and Blood Institute**

National Blood Resources Education Program  
Chairman, Data Safety and Monitoring Committee,  
TRAP Study, 1991-6  
Chairman, Data Safety & Monitoring Committee, T-Cell  
Depletion Trial, 1994-2002

**National Institute on Aging**

Clinical Research Subpanel, 1982-5

**National Institute of Allergy and Infectious Diseases**

Expert Panel on the Effects of Donor Pool Size on the Safety and Efficacy  
of Immunoglobulin Products, 1998

**U.S. Public Health Service**

Advisory Committee on Blood Safety and Availability, 2001- present  
PHS Executive Task Force on AIDS, Charlottesville, VA, 1988  
Medical Review Board, 1982-96  
USPHS Working Group on HTLV-1, 1988  
Medical Board, Clinical Center, NIH, 1982-4, 1989-91  
NIH Strategic Planning Committee (Molecular Medicine  
Initiative Subcommittee), 1992  
Secretary's Committee on Clinical Center Options, 1995

### **U.S. Pharmacopeia**

Committee of Revision, Chairman: Blood and Blood  
Products, 1995-2000  
Committee of Experts – Chairman, Blood & Blood Product Standards,  
2000-2005

### **World Apheresis Association**

Board of Directors, 1993-5  
Councillor - North America, 1986-88, 1992-4  
Scientific Program Committee, 1984, 1986, 1992, 1994

### **World Health Organization**

Consultant to Global Blood Program - 2001

Invited lecturer (including plenary or keynote lectures) at numerous national and international meetings.

Ad hoc reviewer for numerous scientific and medical journals.

## **BIBLIOGRAPHY:**

### **PEER REVIEWED JOURNALS:**

Klein HG, Bell W. Disseminated Intravascular Coagulation Occurring During Heparin Therapy. *Ann Int Med* 1974; 80:477-481.

Klein HG, Aledort LM, Bouma BN., et al. A Cooperative Study for the Detection of the Carrier State of Classic Hemophilia. *N Engl J Med* 1977; 296:959-962.

Klein HG. A Chimpanzee Breeding Colony for Hepatitis Research: *J Med Primat* 1978; 7:182-184.

Weiss GB, Nienhuis AW, McIntosh CL, Klein HG. Traumatic Cardiac Hemolytic Anemia as a Late complication of a Starr-Edwards Mital Valve Prosthesis. *Arch Intern Med* 1979; 139:373-375.

Klein HG, Faltz LA, McIntosh CL, Appelbaum F, Deisseroth AB, Holland PV. Surgical Hypothermia in a Patient with a Cold Agglutinin: Management by Plasma Exchange. *Transfusion* 1980; 20:354-357.

Miller DM, Winslow RM, Klein HG, Wilson KC, Statham NJ, Brown FL, Fulmer J. Improved Exercise Performance after Exchange Transfusion in Subjects with Sick Cell Anemia. *Blood* 1980;56:1127-32.

Klein HG, Garner RJ, Miller DM, Rosen S, Statham NJ, Winslow RM. Automated Partial Exchange Transfusion in Sick Cell Anemia. *Transfusion*, 1980; 20:578-84.

Sazama K, Klein HG, Davey RJ, Corash L. Intraoperative Hemolysis as the Initial Manifestation of Glucose-6-Phosphate Dehydrogenase Deficiency. *Arch Intern Med* 1980; 140:845-6.

Kolins J, Allgood JW, Burghardt D, Klein HG, McGinniss MH. Modification of B, I, i, and Lewis Antigens in a patient with DiGuglielmo's erythroleukemia. *Transfusion* 1980; 20:574-77.

Diamond WJ, Brown F, Bitterman P, Klein HG, Davey R, Winslow RM. Delayed Hemolytic Transfusion Reaction Presenting as Sick Cell Crisis. *Ann Int Med* 1980; 93:231-34.

- Coles SM, Klein HG, Holland PV. Alloimmunization in two multi-transfused patient populations: Sick Cell Disease and Thalassemia Major. *Transfusion* 1981; 21:462-67.
- Corash L, Klein HG, Deisseroth A, Shafer B, Rosen S, Beaman J, Griffith P, Nienhuis A. Selective Isolation of Young Erythrocytes for Transfusion Support of Thalassemia Major Patients. *Blood* 1981; 57:599-606.
- Gadek JE, Klein HG, Holland PV, Crystal RG. Replacement Therapy of Alpha 1-antitrypsin Deficiency: Reversal of Protease-Imbalance within the Alveolar Structures of Piz Subjects. *J Clin Invest* 1981; 68:1158-65.
- Kessler CV, Klein HG, Havlik RJ. Uncontrolled Thrombocythemia in Chronic Myeloproliferative Disorders. *Brit J Haem* 1982; 50(1):157-67.
- Klein HG, Kessler C, Anderson KC, Loomis KF. Multiple Arterial and Venous Thromboemboli in a Male with Chronic Myelogenous Leukemia and Occult Metastatic Breast Carcinoma. *South Med J*. 1982; 75:745-747.
- Lenes B, Klein HG, Lakatos E. Selective Removal of Sick Cells with the IBM 2997 Continuous Flow Blood Cell Separator. *J. Clin. Apheresis* 1983; 1:64-70.
- Young NS, Klein HG, Griffith P, Nienhuis AW. A trial of immunotherapy in aplastic anemia and pure red cell aplasia. *J. Clin. Apheresis* 1983; 1:95-103.
- Reichert CM, Weisenthal LM, Klein HG. Delayed Hemorrhage Following Percutaneous Liver Biopsy. *J. Clin. Gastroenterol.* 1983; 5:263-266.
- Bracey A, Klein HG, Chambers S, Corash L. Simplified Selective Isolation of Young Red Cells for Chronic Transfusion Support. *Blood* 1983; 61:1068-1071.
- Winslow RM, Butler WM, Kark JA, Klein HG, Moo-Penn W. The effects of bloodletting on exercise performance in a subject with a high-affinity hemoglobin variant. *Blood* 1983; 62: 1159-1164.
- Klein HG, McCurdy PR, Braine HG, MacPherson JL, Dowling RN. Centrifugal Leukocytapheresis and Thrombocytapheresis: Complications Related to Equipment and Technique. *J. Clin. Apheresis* 1983; 1:190-195.
- Rose J, Klein H, Greenstein J, McFarlin D, Gerber L, McFarland H. Lymphocytapheresis in chronic progressive multiple sclerosis: Results of a preliminary trial. *Ann Neurol* 1983; 14:593-594.
- Marti GE, Folks T, Longo DL, Klein HG. Concise review: Therapeutic Cytophoresis in chronic lymphocytic leukemia. *J Clin Apheresis* 1983; 1:243-248.

Dowling R, Weber V, Osborne L, Klein HG. Mononuclear cell collection using various techniques. *J Clin Apheresis* 1984; 2:32-40.

Lane HC, Masur H, Longo DL, Klein HG, Rook AH, Quinnan GV, Steis RG, Macher A, Whalen G, Edgar LC, Fauci AS. Partial Immune Reconstitution in a Patient with the Acquired Immunodeficiency Syndrome. *N Engl J Med* 1984; 311: 1099-1109.

Rodgers GP, Bonner RF, Klein HG, Noguchi CT, Nienhuis AW, Schechter AN. Periodic Microcirculatory Flow in Patients with Sickle Cell Disease. *N Engl J Med* 1984; 311: 1534-8.

Grindon AJ, Tomasulo PA, Bergin JJ, Klein HG, Miller JD, Mintz PD. The Hospital Transfusion Committee: Guidelines for Improving Services. *JAMA* 1985; 253:540-543.

Winslow RM, Klein HG, Monge CC. Red cell mass in Andean natives with excessive polycythemia. *Arch Biol Andina* 13:85-94, 1984-85.

Winslow RM, Monge CC, Brown EG, Klein HG, Sarnquist F, Winslow NJ. The effect of hemodilution on O<sub>2</sub> transport in high altitude polycythemia. *J Appl Physiol* 1985; 59(5): 1495-1502.

Tomasulo PA, Lenes BA, Noto TA, Klein HG, Menitove J. Automatic Special Case Consultations in Transfusion Medicine. *Transfusion*, 1986; 26:186-193.

Klein HG, Balow JE, Dau PC, Hamburger MI, Leitman SF, Pineda AA, and Tindal RSA: Clinical Application of Therapeutic Apheresis. *J Clin Apheresis* 1986; 3(7):1-93.

AuBuchon JP, Carter CS, Adde MA, Meyer DR, Klein HG. Optimization of Parameters for Maximization of Plateletpheresis and Lymphocytophoresis yields on the Haemonetics Model V50. *J. Clin. Apheresis* 1986; 3(2):103-110.

Stevenson HC, Stevenson GW, Leitman SF, Carter CS, Alvord G, Klein HG. The isolation of human mononuclear subsets by cytophoresis for laboratory research. *Plasma Ther Transfus Technol* 1986; 7:365-371.

Muul LM, Nason-Burchenal K, Carter CS, Cullis H, Slavin D, Hyatt C, Director EP, Leitman SF, Klein HG, Rosenberg SA. Development of an automated closed system for generation of human lymphokine activated killer (LAK) cells for use in adoptive immunotherapy. *J Immunol Methods* 1987; 101:171-181.

Carter CS, Leitman SF, Cullis H, Muul LM, Nason - Burchenal K, Rosenberg SA, Klein HG. Use of a continuous flow cell separator in density gradient isolation of lymphocytes. *Transfusion* 1987; 27:362-365.

- Areman EM, Simonis T, Carter C, Read EJ, Klein HG. Bulk cryopreservation of lymphocytes in glycerol. *Transfusion* 1988; 28:151-156.
- Rodgers GP, Schechter AN, Noguchi CT, Klein HG, Nienhuis AW, Bonner RF. Microcirculatory adaptations in sickle cell anemia. Reactive hyperemia response. *Am J Physiol* 1990;258:
- Kruskall MS, Mintz PD, Bergin JJ, Johnston MFM, Klein HG, Miller JD, Rutman R, Silberstein L. Transfusion therapy in emergency medicine. *Ann Emergency Med* 1988; 17:327-335.
- Sloand EM, Kessler CM, McIntosh CL, Klein HG. Neutralization of heparin-induced anticoagulation with methylene blue. *Thromb. Res.* 1989, 54:677-86.
- Carter CS, Leitman SF, Cullis H, Muul LM, Nason-Burchenal K, Rosenberg SA, Klein HG. Technical Aspects of lymphokine-activated killer cell production. *J. Clin. Apheresis* 1988; 2/3:113-117.
- Leitman SF, Klein HG, Melpolder JJ, Shih JW, Read EJ, Harvarth L, Darr F, Foy JL, Alter HJ. Clinical implications of positive tests for antibodies to human immunodeficiency virus type 1 in asymptomatic blood donors. *N Engl J Med* 1989; 321:917-23.
- Sloand EM, Fox SM, Banks SM, Klein HG. Preparation of IgA deficient platelets. *Transfusion* 1990; 30:322-326.
- Loiacono B, Carter G, Leitman S, Klein HG. Efficacy of various methods of confidential unit exclusion in identifying potentially infectious blood donations. *Transfusion* 1989; 29:823-6.
- DePalma L, Palmer R, Leitman SF, Dolan WD, Klein HG. Utilization patterns of frozen autologous red blood cells: experience in a referral center and a community hospital. *Arch Pathol Lab Med* 1990; 114:516-18.
- Bergsten P, Amitai G, Kehrl J, Dhariwal KR, Klein HG, Levine M. Millimolar concentrations of ascorbic acid in purified human mononuclear leukocytes: depletion and reaccumulation. *J Biol Chem* 1990; 265:2584-2587.
- Sloand EM, Klein HG. Effect of white cells on platelets during storage. *Transfusion* 1990; 30:333-338.
- Lane HC, Zurich KM, Wilson W, Cefali F, Easter M, Kovacs JA, Masur H, Leitman SF, Klein HG, Steis RG, Longo DL, Fauci AS. Syngeneic bone marrow transplantation and adoptive transfer of peripheral blood lymphocytes combined with zidovudine in human immunodeficiency virus infection. *Ann Int Med* 1990; 113:512-19.

Alter HJ, Epstein JS, Swenson SG, Van Raden MA, Klein HG, Ward JW, Kaslow RA, Menitove JE, Sandler SG, Sayers MH, Hewlett IK, Chernoff AI. Prevalence of Human Immunodeficiency Virus Type 1 (HIV-1) p24 antigen in U.S. Blood Donor: Assessment of Efficacy for Blood Donor Screening. *N Engl J Med* 1990; 323:1312-17.

Sloand EM, Sloand JA, Prodouz K, Klein HG, Yu MW, Harvath L, Fricke W. Reduction of platelet glycoprotein Ib in uremia. *Brit J. Haem* 1991; 77:375-81.

Fricke W, Alling D, Kimball J, Griffith P, Klein H. Lack of efficacy of tranexamic acid in thrombocytopenic bleeding. *Transfusion* 1991; 31:345-348.

Keilholz U, Klein HG, Korbling M, Brado B, Carter CS, Cullis H, Galin F, Hunstein W. Peripheral blood mononuclear cells collection from patients undergoing adoptive immunotherapy or peripheral blood-derived stem cell transplantation and from normal donors. *J. Clin Apheresis* 1991; 6:131-6.

Sloand EM, Klein HG, Pastakia KP, Pierce P, Prodouz KN. Effects of albumin on the inhibition of platelet aggregation by beta lactam antibiotics. *Blood* 1992; 79:2022-27.

Sloand EM, Klein HG, Banks SM, Vareldzis B, Merritt S, Pierce P. Epidemiology of thrombocytopenia in HIV infection. *Eur J. Haem* 1992; 48:168-172.

Natanson C, Hoffman WD, Koev LA, Dolan DP, Banks SM, Bacher J, Danner RL, Klein HG, Parrillo JE. Plasma exchange does not improve survival in a canine model of human septic shock. *Transfusion* 1993; 33:243-8.

Stroncek DF, Holland PV, Bartsch G, Bixby T, Simmons RG, Antin JH, Anderson KC, ASH PC, Bolwell BH, Hansen JH, Heal JM, Henslee-Downey PJ, Jaffe' ER, Klein HG, Lau PM, Perkins HA, Popovsky MA, Price TH, Rowley SD, Stehling LC, Weiden PL, Wissel ME, McCullough J. Experiences of the first 493 unrelated marrow donors in the National Marrow Donor Program. *Blood* 1993; 81:1940-46.

Sloand E, Kumar P, Klein HG, Merritt S, Sacher R. Transfusion of blood products in HIV-1 infected persons: relationship to opportunistic infection. *Transfusion* 1994; 34:48-53.

Sloand EM, Kumar P, Yu M, Klein HG. Effect of amphotericin B and fluconazole on platelet membrane glycoproteins. *Transfusion* 1994; 34: 415-20.

Sloand EM, Alyono D, Klein HG, Chang P, Yu M, Lightfoot FG, Kessler C. 1-Deamino-8-D-arginine vasopressin (DDAVP) increases platelet membrane expression of glycoprotein 1b in patients with disorders of platelet function and after cardiopulmonary bypass. *Am J. Hematol* 1994; 46: 199-207.

Lubarsky DA, Hahn C, Bennett DH, Smith LR, Bredehoeft SJ, Klein HG, Reves DG. The hospital costs (fiscal year 1991/1992) of a simple perioperative allogeneic red blood cell transfusion at Duke University. *Anesth Analg* 1994; 79:629-37.

Lee JH, Klein HG. The effect of donor erythrocyte sedimentation rate on efficiency of granulocyte collection by centrifugal leukapheresis. *Transfusion* 1995; 35:384-388.

Lee JH, Cullis H, Leitman SF, Klein HG. The efficacy of pentastarch in granulocyte collection by centrifugal leukapheresis. *J Clin Apheresis*. 1995;10:198-202.

Lee JH, Leitman SF, Klein HG. A controlled comparison of the efficacy of hetastarch and pentastarch in granulocyte collections of centrifugal leukapheresis. *Blood* 1995; 86:4662-6.

Sloand EM, Alyono D, Yu M, Klein HG. Platelet membrane glycoproteins and microvesicles in blood from post-operative salvage: a study in cardiac bypass patients. *Transfusion* 1995; 35:738-744.

Blaese RM, Culver KW, Miller AD, Carter CS, Fleisher T, Clerici M, Shearer G, Chang L, Chiang Y, Tolstosher P, Greenblatt J, Rosenberg S, Klein HG, Berger M, Mullen C, Ramsey J, Muul L, Morgan RA, Anderson WF. T lymphocyte-directed gene therapy for ADA Deficiency (SCID): Initial trial results after 4 years. 1995; *Science* 270:475.

Sloand EM, Yu M, Klein HG. Comparison of random-donor platelets prepared from whole blood units and single-donor platelet's prepared from single donor apheresis collections. *Transfusion* 1996; 36:955.

Read EJ, Kunitake ST, Carter CS, Chau Q, Klein HG. Enumeration of CD34+ hematopoietic progenitor cells in peripheral blood and leukapheresis products by microvolume fluorimetry: a comparison with flow cytometry. *J Hematotherapy*. 1997, 6:291-301.

Walker RE, Carter CS, Muul L, Natasajan V, Herpin BR, Leitman SF, Klein HG, Mullen CA, Baseler M, Falloon J, Davey Jr RT, Kovacs JA, Polis MA, Masur H, Blaese RM, Lane HC. Peripheral expansion of mature T Cells is a major means of CD4+ T cell regeneration in adults with HIV-1 infection. *Nature Med*. 1998, 4:852-6.

Silva N, O'Bryan L, Medeiros E, Holand H, Suliman J, Patronas N, Reed SG, Klein HG, Masur H, Badaro R, Trypanosoma Cruzi meningoencephalitis in HIV-infected patients. *J. Acquir Immun Defic Syndr Hum Retrovirol* 1999; 20:342-9.

Zumberg MS, Procter JL, Lottenberg R, Kitchens CS, Klein HG. Autoantibody formation in the alloimmunized red blood cell recipient. *Arch Int Med* 2001; 161:281-90.



Stroncek DF, Procter JL, Moses L, Bolan C, Pomper GJ, Conry-Cantilena C, Malech HL, Klein, HG, Leitman, SF. Intravenous Rh immune globin prevents alloimmunization in D negative granulocyte recipients, but obscures the detection of an alloanti-K. *Immunology* 2001, 17:37-41.

Wang E, Adams S, Panelli M, Klein H, Marincola FM. Fluorimetric detection of single nucleotide variants (SNV) by proportional hybridization to oligonucleotide arrays (PHOA). *Nature Biotechnology* (in press)

Stroncek DF, Rainer T, Sharon V, et al. Sick hemoglobin polymerization in red cell components from donors with sickle trait prevents effective leukocyte reduction by filtration. *Transfusion* 2002;42:1466-72

Panelli MC, Wang E, Phan G, Puhlman M, Miller L, Ohnmacht GA, Klein HG, Marincola FM. Gene expression profiling of the response of peripheral blood mononuclear cells and melanoma metastases to systemic IL-2 administration. *Genome Biol* 2002; 25:3(7) Research 0035

Haddad SA, Winer KK, Gupta A, Chakrabarti S, Noel P, Klein HG. A puzzling case of anemia. *Transfusion* 2002;42:1610-13.

Wang E, Adams S, Zhao Y, Panelli MC, Simon R, Klein H and Marincola FM. A strategy for detection of known and unknown SNP using a minimum number of oligonucleotides applicable in the clinical settings *Journal of Translational Medicine* 2003, 1:4 (online; 20 Aug 2003)

Leitman SF, Browning JM, Yau YY, Mason G, Klein HG, Conry-Cantilena C, Bolan CD. Impact of using hemochromatosis subjects as allogeneic blood donors: a prospective study. *Transfusion* (in press)

Guralnick J, Woodman R, Klein HG et al. Anemia and functional status in an elderly population. *JAMA* (in press)

## **REVIEWS:**

Borberg H, Dahlke MB, Fraser ID, Fratantoni JC, French JE, Holland PV, Klein HG, Polesky HF, Sandler SG. Which are the established or potential risks for Donors undergoing Cytapheresis Procedures and how can they be prevented? *Vox Sang* 1980; 39(3):169-79.

Klein HG. Selective Isolation of Young Red Cells for Chronic Transfusion. *Plasma Therapy* 1981; 2:175-80.

Klein HG, Winslow RW. Partial Exchange Transfusion in Sick Cell Disease. *Prog Clin Biol Res* 1982; 88:235-243.

Klein HG. Transfusions with young erythrocytes (neocytes) in Sick Cell Anemia. *Am J Ped Hemat Onc* 1982; 4(2):162-165.

Klein HG. Therapeutic Cytapheresis. *Prog Clin Biol Res* 1982; 106:297-306.

Klein HG. Therapeutic Apheresis: An overview. In: Nose, Y., Malchesky, P.S., and Smith, J.W. (Eds). *Plasmapheresis* ISAO Press (Cleveland) 1983:3-6.

Klein HG. Therapeutic Cytapheresis. In: Kolins J, Jones J, eds. *Therapeutic Apheresis*, Washington: American Association of Blood Banks, 1983:23-34.

Klein HG. The Kinship of Coagulation and Immunity: The Contact System. In Tregellas WM and Keating LJ, eds. *Immunology*. Arlington, VA: American Association of Blood Banks, 1985:177-190.

Klein HG, Alter HJ. Blood Transfusion and AIDS In: *AIDS*, Chicago: American Medical Association 1987:7-10.

Sloand EM and Klein HG. Therapeutic Apheresis. In: Kasprisin DO (ed) *Current Status of Hemapheresis*, Washington: American Association of Blood Banks 1987:9-47.

Klein HG, Leitman SF. Adoptive immunotherapy in malignant disease. *Transfusion* 1989; 29:170-178.

Higgins MJ, Klein HG. Massive transfusion in the intensive care unit. *J Intensive Care Med* 1989; 4:221-233.

The use of autologous blood. The National Blood Resource Education Program Expert Panel. *JAMA* 1990; 263:414-17.

Klein HG. Controversies in transfusion medicine: Alanine aminotransferase screening of blood donors: Pro. *Transfusion* 1990; 30:363-367.

Klein HG. When is Transfusion the best option? *Emergency Medicine*. 1992; 24(5): 59-66.

Strauss RG, Ciavarella D, Gilcher RO, Kasprisin DO, Kiproff DD, Klein HG, McLeod BC. Clinical Application of Therapeutic Hemapheresis. *J. Clin Apheresis* 1993; 8:189-269.

Walker R, Blaese RM, Carter CS, Chang L, Klein H, Lane HC, Leitman SF, Mullen CA, Larson M. A study of the safety and survival of the adoptive transfer of genetically marked syngeneic lymphocytes in HIV-1 infected identical twins. *Human Gene Ther* 4 (5) 1993; 659-80.

Klein HG. Immunologic Aspects of Blood Transfusion. Seminars in Oncology 1994; 21:16-20.

Lee JH, Klein H.G. Mononuclear cell adoptive immunotherapy Hematology/Oncology Clinics of North America 1994; 8(6):1203-1221.

Lee JH, Klein HG. Peripheral blood stem cell transplantation Hematology/Oncology Clinics of North America 1994; 9(1):1-22.

Lee JH, Klein HG. Cellular gene therapy. Hematology/Oncology Clinics of North America 1994; 9(1):91-113.

Sloand EM, Pitt E, Klein HG. Safety of the blood supply. JAMA 1995; 274:1368-73.

Klein HG. Allogeneic transfusion risks in surgical patients. Am J. Surg 1995; 170 (suppl): 215-265.

Hillyer CD, Klein HG. Immunotherapy and gene transfer in the treatment of the oncology patient: Role of transfusion medicine. Transf Med Rev 1996; 10(1):1-8.

Miller Y, Klein HG. Growth factors and their impact on Transfusion Medicine. Vox Sang 1996; 71:196-204.

Klein HG. New insights into the management of anemia in the surgical patient. Am J Med 1996; 101 (2A): 125-155.

Klein HG, Strauss RG, Schiffer CA. Granulocyte transfusion therapy. Semin Hematol 1996; 33:359-368.

Klein HG, Dodd RY, Dzik WH, Luban NLC, Ness PM, Pisciotto P, Schiff P, Snyder EL. Current status of solvent/detergent-treated frozen plasma. Transfusion 1998; 38:102-107.

Klein HG. Immunologic Aspects of Transfusion: A once and future risk? Anesthesiology 1999; 91:861-5.

Engelfriet CP, Reesink HW, Blajchman MA et al. Bacterial contamination of blood components. Vox Sanguinis 2000; 78:59-67.

Engelfriet CP, Reesink HW, Klein HG, et.al. International Forum: granulocyte concentrates. Vox Sang 2000; 79:59-66.

Klein HG. The immunomodulatory effects of blood transfusion. Tumori 2001; 87(2) 18-20

Engelfriet CP, Reesink HW, Klein H.G. et al. The future of pathogen-inactivated platelet concentrates. *Vox Sang* 2003; 85:54-66.

Brittenham GM, Klein HG, Kushner JP, Ajioka RS. Preserving the national blood supply. *Hematology (Am Soc Hematol Educ Program)*. 2001; 422-32

#### **BOOK CHAPTERS:**

Klein HG. Effects of plasma exchange on plasma constituents: Choice of replacement solutions. In: MacPherson JL and Kasprisin DO (Eds). *Therapeutic Hemapheresis* CRC Press Inc. (Boca Raton) Vol II, 1985:3-14.

Klein HG. Erythrocytapheresis. In: Valbonesi M, Pineda AA, Biggs JC, eds. *Therapeutic Hemapheresis*, Milan: Wichtig Editore. 1986:143-148.

Read EJ, Klein HG. Hematological Effects of Aging: Considerations for Clinical Trials. Cutler NR, Narang PK, eds. *Drug Studies in the Elderly: Methodological Concerns*, New York: Plenum 1986:123-150.

Klein HG. Blood Product Therapy in the Critical Care Unit. Parrillo JE, ed. *Current Therapy in Critical Care Medicine*, Philadelphia: B.C. Decker Inc. 1987:281-287.

Klein HG. Transfusion support of hemoglobinopathies. In: Cash J (ed) *Progress in Transfusion Medicine* Vol III. Churchill Livingstone (Edinburgh) 1987: 198-213.

Pineda AA, Klein HG. Applications of Hemapheresis. In: Greenwalt TJ (Ed). *Methods in Hematology*, Vol 17. Blood Transfusion, New York: Churchill Livingstone (Edinburgh). 1988: 213-233.

Klein HG. Young red cells for transfusion support. In: Pineda AA, Valbonesi M, Sniecinski IJ, eds. *Donor Hemapheresis*, Milan, Italy: Wichtig Editore, 1988: p. 65-69.

Lacerna LV, Hester J, Pineda AA, Burgstaler E, Klein HG. Clinical/Technical challenges in adoptive cellular immunotherapy: The role of cytapheresis, In: Stevenson HC (Ed) *Adoptive Cellular Immunotherapy of Cancer*, New York: Marcel Dekker, Inc. 1989: 175-190.

Klein HG, Travis LB. Transfusion support of hemoglobinopathies. In: Dutcher JP (ed) *Modern Transfusion Therapy*, Vol I, CRC Press Inc., Boca Raton, 1990: p. 93-106.

Klein HG. Apheresis. In: Hoffman R, Benz EJ, Shattil S, Furie B, Cohen HJ, eds. *Hematology: Principles and Practice*. New York: Churchill Livingstone Inc. 1990: p1638-43.

Klein HG, Apheresis in transition. In: Nydegger UE, (ed): Therapeutic Hemapheresis in the 1990's. Curr Stud Hematol Blood Transf. Basel, Karger, 1990, No. 57, pp 1-9.

Klein HG. Plasma exchange in the future: innovations and new indications. In Rossi EC, Simon TL and Moss GS (eds): Principles of Transfusion Medicine. Williams and Wilkins, Baltimore 1991: p557-560.

Klein HG. Therapeutic mononuclear cell transfusion: adoptive immunotherapy. In Rossi EC, Simon TL, and Moss GS (eds): Principles of Transfusion Medicine, Williams and Wilkins, Baltimore. 1991: p295-300.

Klein HG. Autologous mononuclear cell therapy. In Taswell HF and Pineda AA (eds): Autologous Transfusion and Hemotherapy. Blackwell Scientific Publication, Boston. 1991: p220-232.

Leonard K, Klein HG. Acute Hemolytic Anemia. In Bell WR (ed): Hematologic and Oncologic Emergencies. Contemporary Management in Critical Care. Churchill Livingstone, New York 1993: pp33-69.

Klein, H.G. Immunohematology and Blood Transfusion. In Isselbacher KJ, Braunwald E, Wilson JD, Martin JB, Fauci AS, Kasper DL (eds): Harrison's Principles of Internal Medicine, McGraw-Hill Inc, New York 1993: p 2377.

Klein HG. Principles of Apheresis. In Anderson KC and Ness PM (eds). Scientific Basis of Transfusion Medicine. WB Saunders, Philadelphia 1994: p 781-796.

Cantilena CC, Klein HG. Therapeutic use of blood components. In Rakel CT (ed): Conn's Current Therapy. W.B. Saunders Company, Philadelphia, PA 1994: p 427-31.

Higgins MJ, Klein HG. Blood Products in the ICU. In Parrillo JE and Bone RC (eds): Critical Care Medicine: Principles of Diagnosis and Management. Mosby-Year Book, Inc. Philadelphia, PA 1995, p 1323-44.

Klein HG. Hemapheresis and Cellular Therapy. In Hoffman R, Benz EJ, Shattil S, Furie B, Cohen HJ (eds): Hematology: Principles and Practice, New York: Churchill Livingstone Inc. 1995: p2038-2045.

Klein HG. Therapeutic Hemapheresis. In: Petz LD, Swisher S, Kleinman S, Spence R, Strauss RG, (eds): Clinical Practices of Transfusion Medicine. Churchill Livingstone, New York 1996, p1011-1022.

Carter CS, Gress R, Klein HG. Routine methods for preparing hematopoietic progenitor cell components. In Lasky L and Warkentin P (eds): Marrow and Skin Cell Processing for Transplantation. American Association of Blood Banks, Bethesda, MD 1995, p 51-68.

Klein HG. Immunohematology, In: Rich RR, Fleisher TA, Schwartz BD, Shearer WT, Strober W (eds): Clinical Immunology: Principles and Practice. Mosby, St. Louis 1995, p2220-2228.

Klein HG. Novel Cellular Therapies, In: Van Oss CJ (ed). Transfusion Immunology and Medicine. Marcel Dekker, Inc. (New York) 1995; 411.

Klein HG. The immunomodulatory effects of blood transfusion. In: Petz LD, Swisher S, Kleinman S, Spence R, Strauss RG (eds): Clinical Practice of Transfusion Medicine. Churchill Livingstone, New York 1996, p 59-70.

Klein HG. Therapeutic Hemapheresis. In: Petz LD, Swisher S, Kleinman S, Spence R, Strauss RG, (eds) : Clinical Practices of Transfusion Medicine. Churchill Livingston, New York 1996, p 1011-1022.

Lee JH, Klein HG. Therapeutic mononuclear cell transfusion: adoptive immunotherapy and gene therapy. In: Rosse EC, Simon TL, and Moss GS (eds): Principles of Transfusion Medicine. Williams and Wilkins, Baltimore 1996, p 329-44.

Higgins MS, Klein HG. Blood Products in the Critical Care Unit. Parrillo JE ed. Current Therapy in Critical Care Medicine. Mosby-Year Books, Inc. Philadelphia 1997, p 362-7.

Klein HG. Cellular Gene Therapy. In McLeod B, Price TH, Drew MJ (eds): Principles of Apheresis. AABB Press, 1997, p 537-46 .

Klein HG. Continuing issues regarding transfusion and coagulation factor-acquired HIV infection. In: Pizzo PA, Wilfert CM (eds); Pediatric AIDS. (3<sup>rd</sup> Edition) Baltimore, Williams and Wilkins, 1998.

Lee JH, Klein HG. Mononuclear cell transfusion: Immunotherapy Using Allogeneic Donor-Derived Lymphocytes. Mintz PD (ed) Transfusion Therapy: Clinical Principles and Practice. AABB Press, Bethesda 1998, p 251-265.

Klein HG. Principles of Apheresis. In: Anderson KC and Ness PM (eds) Scientific Basis of Transfusion Medicine. WB Saunders, Philadelphia 1999: p553-68.

Conry-Cantilena, Klein, HG. Hemochromatosis subjects as blood and tissue donors. In: Barton JC and Edwards CO, (eds); Hemochromatosis. Cambridge University Press, Cambridge 2000: p564-73.

Lazarus EF, Klein HG. Hemapheresis and cellular therapy. In: Hoffman R. Benz EJ, Shattil S, Furie B, Cohen HJ, Silberstein LE, McGlave P (eds): Hematology: Principles and Practice. Churchill Livingstone Inc.: New York 1999, p2291-99.

Klein HG. Molecular Biology in Transfusion Medicine. In: Smit Sibinga C Th and Klein HG (eds). Molecular Biology in Transfusion Medicine. Kluwer Academic Publishers (Dordrecht) 2000, p3-12..

Kicklighter EJ, Klein HG. Hemolytic transfusion reactions. In: Linden JV, Bianco (eds). Blood Safety and Surveillance, Marcel Dekker, inc., New York 2001, p47-70..

Lee, J-H, Klein HG. From leukocyte reduction to leukocyte transfusion: the immunologic effects of transfused leukocytes. Contreras M (ed) Bailliere Tindal, London 2000, p585-600.

Griffith L, Klein HG. Immunohematology, In: Rich RR, Fleisher TA, Kotzin B, Shearer WT, Schroeder H (eds): Clinical Immunology: Principles and Practice. Churchill Livingstone, London 2001, p127.1-127.10.

Lazarus EF, Klein HG. Apheresis, In Rich RR, Fleisher TA, Kotzin B, Shearer WT, Schroeder H (eds): Clinical Immunology: Principles and Practices. Churchill Livingstone, London, 2001, p118.1-118.9

Bolan CD, Klein HG. Transfusion Medicine and pharmacologic aspects of hemostasis. In Kitchens CS, Alving BM, Kessler CM (eds): Consultative Hemostasis and Thrombosis. W.B. Saunders Company, New York 2002, 395-418.

Klein HG. The evolution of cellular therapy in the new millennium. In: Sacher RA (ed) Cellular Therapy: New Frontiers in transfusion Medicine. AABB Pres (Bethesda) 2002, 1-13.

Klein HG. Cellular gene therapy. In: McLeod BC (ed) Apheresis: Principles and Practice. AABB Press (Bethesda) In Press

Lazarus EF, Klein HG. Hemapheresis and cellular therapy. In: Hoffman R, Benz EJ, Shattil S, Furie B, Cohen HJ, Silberstein LE, McGlave P (eds): Hematology: Principles and Practice. Churchill Livingstone Inc.: New York (in press)

## **GOVERNMENT PUBLICATIONS:**

Petit CR, Klein HG. Hemophilia, Hemophiliacs and the Health Care System. DHEW Publ. No. (NIH) 1975:76-781.

Klein HG. Isovolemic Hemodilution in High Altitude Polycythemia. In: Adjustment to High Altitude, DHHS Publ. No. (NIH) 1983; 83-2496: 47-52.

Anderson D, Epstein J, Pierik L, Solomon J, Blattner W, Saxinger L, Alter H, Klein H, McCurdy P, Nemo G, Kaplan J, Allen J, Khabbaz R, Lairmore M. Licensure of screening tests for antibody against human T-lymphocytotropic virus type I. MMWR 37(48):736-740, December 9, 1988.

Toy PT, Dzik W, Gould SA, Hoppe PA, Kruskall MA, Maurer RC, Menitove JE, Naiman JL, Thurer RL, Klein HG, McCurdy PR, Sloand EM. Use of Autologous Blood. U.S. Department of Health and Human Services Publ. No. (NIH) 89-3038.

Stehling LC, Cosgrove DM, Moss GS, Schiffer CA, Slichter SJ, Snyder EJ, Klein HG, McCurdy PR, Sloand EM. Indications for the use of red blood cells, platelets, and fresh frozen plasma. U.S. Department of Health and Human Services. Publ. No. (NIH) 89-2974.

Toy PT, Beattie C, Gould SA, et al. Transfusion Alert. Use of Autologous Blood. U.S. Department of Health and Human Services Publ. No. (NIH) 94-3038.

#### **PUBLISHED PROCEEDINGS:**

Klein HG, Alter HJ, Holland PV. Postexposure immunoglobulin prophylaxis of hepatitis B: A comparison of two dosage schedules. In Vyas GN, Cohen SM, Schmid R (Ed): Viral Hepatitis: Proceedings of the Second Symposium of Viral Hepatitis (Franklin Institute Press, Philadelphia) 1978:483-86.

Klein HG. Neocytapheresis: Collection, Preparation and Preliminary Clinical Results, in Proceedings of the Red Cross Apheresis Symposium, Chicago, American Red Cross Blood Services, Mid America Region. 1979:88-96.

Klein HG. Red Cell Exchange Transfusion in Secondary Polycythemia. In Proceedings of the Red Cross Second Apheresis Symposium, Chicago, 1980:74-82.

Spiro T, Young N, Klein HG. Collection of Erythroid (BFU-e) and Myeloid (CFU-c) Progenitor Cells from Peripheral Blood using Continuous Flow and Intermittent Flow Cell Separators. in Proceedings of the Red Cross Second Apheresis Symposium, Chicago, 1980; 158-72.

Klein HG. Cell Separators for Red Cell Exchange in Advances in the Pathophysiology, Diagnosis, and Treatment of Sickle Cell Disease (Alan R. Liss, Inc., New York) 1982; 109-117.

Schulz SC, van Kammen DP, Waters R, Klein HG, Balow JE, Bunney WE. Double Blind Evaluation of Plasmapheresis in Schizophrenic Patients: A Pilot Study. Artificial Organs 1983; 12:3141.



- Young NS, Klein HG, Griffith P, Nienhuis AW. Therapeutic plasma exchange and lymphocyte depletion in aplastic anemia and pure red cell aplasia. In: Nose, Y., Malchesky, P.S., J.W., and Krakauer, R.S. (Eds). Plasmapheresis Raven Press (New York) 1983:339-348.
- Funck-Brentano JL, Klein HG, Lysaght MJ, Oda T. Plasma Exchange or Plasma Treatment. In: Nose Y., Malchesky A.S. and Smith J.W. (eds). Plasmapheresis: New Trends in Therapeutic Applications. ISAO Press (Cleveland) 1983: 167-171.
- Groopman JE, Klein HG, Levine PH, Sande MA, Redfield RR. AIDS: Questions and Controversies - A Panel Discussion. Sem Hem 1987; 24 (3) Suppl 2: 9-19.
- Klein HG. Adoptive immunotherapy: novel applications of blood cell separators. J. Clin Apheresis 1988; 4:198-202.
- Chernoff AI, Klein HG, Sherman LA (Eds). Research opportunities in transfusion medicine. Transfusion 1989; 29:711-742.
- Stevenson HC, Klein HG. Biologic response modifiers (BRM) chemotherapy and activated cytotoxic leukocyte subsets: strategies for rational design of apheresis-based cancer immunotherapies. Prog Clin Biol Res 1990; 337:559-65.
- Immunosuppressive effect of blood transfusion. Klein HG and Tarrter PI (eds) Dept of Surgery, Mt. Sinai Medical Center (New York) 1991.
- Klein HG. Current risks of blood transfusion. In Friedel N, Hetzer R, Royston D, (eds) Blood Use in Cardiac Surgery. Steinkopff Verlag (Darmstadt) 1991:69-73.
- Klein HG. Future Technologies in Transfusion Medicine. Infusionstherapie 1991; 18 (suppl 1):31-34.
- Shih JW-K, Kung K, Dragon EA, Davies C, Pottathil R, Leitman SF, Klein H, Spadaro J, Melpolder J, Alter HJ. Quantitative analysis of an HIV positive blood donor cohort. In Andrieu JM (ed) Viral Quantitation in HIV Infection. John Libbey Eurotext, (Paris) 1991; 171-176.
- Klein HG. Transfusion Medicine: The impact of biotechnology, growth factors, and bioengineering. In Smit Sibinga C. Th, Das PC, and Cash JD. Transfusion Medicine: Fact and Fiction. Kluwer Academic Publishers (Dordrecht) 1991;153-158.
- Cottler-Fox M, Read EJ, Yu M, Carter C, Klein HG. Mononuclear and CD34 stem cell recovery after automated ficoll processing of marrow or peripheral blood for transplantation. In: Gross S, Gee AP, Worthington-White D eds. Prog Clin Biol Res, New York: Wiley-Liss, 1992; 377:569-73.

Klein HG. Transfusion in Transplant Patients: the Good, the Bad and the Ugly. *Journal of Heart/Lung Transplantation* 1993; 12(S):7-12.

Klein HG. Apheresis, immunoreconstitution and AIDS. In Agishi T (ed) *Therapeutic Plasmapheresis (XII)* VSP Publishers (Tokyo) 1993; 21-23.

Cottler-Fox M, Klein HG. Transfusion support of hematology and oncology patients. The role of hematopoietic growth factors. *Arch Pathol Lab Med* 1984; 118:417-20.

Klein HG. Oxygen Carriers and transfusion medicine. *Art Cells Blood Subs and Immob Biotech* 1994; 22:123-135.

Klein HG. Novel cellular therapies. *Immunol Invest* 1995; 24:411-20.

Klein HG. Cellular Gene Therapy: An overview. *J. Clin Apheresis* 1994; 9:139-41.

Klein HG. Immunologic aspects of blood transfusion. *Semin Oncol* 1994; 21 (2), Suppl 3, 16-20.

Klein HG, McCurdy PR, Nemo G, et al. Transfusion alert: Use of autologous blood. *Transfusion* 1995; 35:703-11.

Klein HG. Examples of risks that we tolerate. In: Bendixen H, Manning FJ, Sparacino L (eds). *Blood and Blood Products: Safety and Risk*. National Academy Press (Washington), 1996, p101-105.

Klein HG, Dodd RY, Ness PM, Fratantoni JA, Nemo GJ. Current status of microbial contamination of blood components: summary of a conference. *Transfusion* 1997; 37:95-101.

#### **BOOKS EDITED:**

*Standards for Blood Banks and Transfusion Services*, 16th Edition, American Association of Blood Banks, Bethesda 1994.

*Standards for Blood Banks and Transfusion Services*, 17th Edition, American Association of Blood Banks, Bethesda, MD, 1996.

*Transfusion Medicine*. *Semin Hematol* (Guest Editor), 1996; 33 (4) : 275-358.

*Molecular Biology in Transfusion Medicine*. Smit Sibinga C Th and Klein HG (eds). Kluwer Academic Publishers (Dordrecht) 2000.

#### **EDITORIALS:**

Klein HG. Therapeutic Plasma Exchange: A Healthy Dose of Skepticism. *Western J Med* 1983;92-94.

Klein HG. Blood Transfusion and Athletics; Games People Play. *New Engl J Med* 1985; 312:854-856.

Klein HG. Transfusion Medicine: The evolution of a new discipline. *JAMA* 1987; 258: 2108-2109.

Klein HG. Developmental Milestones: The Journal at five years. *J Clin Apheresis* 1988, 4:1-2.

Klein HG. Wolf in Wolf's clothing: Is it time to raise the bounty on the passenger lymphocyte? *Blood* 1992; 80:1865-8.

Klein HG. It seemed a shame to throw away the red cells. Selective component collection. *Transfusion* 1993; 33:788-90

Klein HG. Preface. In: Manning FG and Sparacino L (eds): *Blood Donors and the Supply of Blood and Blood Products*. National Academy Press (Washington) 1996, p ix-x.

Klein HG. Introduction: Transfusion Medicine. *Semin Hematol* 1996; 33: 275-6.

Blajchman MA, Klein HG. Looking back in anger: Retrospection in the face of a paradigm shift. *Transf Med Rev*. 1997 ; 11 (1) :1-5.

Klein HG. Cellular Gene Therapy: A Credible Odyssey. *Transfusion*. 1997; 37:675-77.

Klein HG. Blood, Drugs, and the United States Pharmacopeia. *Transf Med Rev* 1998; 12(2):81-3.

Klein HG. Transfusion Safety: Avoiding unnecessary bloodshed. *Proc. Mayo Clin* 2000, 75: 5-7.

Klein HG. The prospect of red-cell substitutes. *N Engl J Med* 2000; 342: 1666-8.

Klein HG. Will blood transfusion ever be safe enough? *JAMA* 2000; 284(2): 238-240.

Klein HG. Blood donation: It's more than comfort and convenience. *AABB News* 2001; January/February: 2.

Klein HG. The transfusion timeline and current good health care practices. *AABB News* 2001; May/June: 2

Klein HG. Safety and availability: The CJD paradigm. AABB News 2001; July/August:2.

Klein HG. Rushing into the new millennium. AABB News 2001; September/October: 1

Klein HG. Earthquake in American. Transfusion 2001. 41:1179-80.

Klein HG. Older is not necessarily better. Anesthesiology 2003 98(4):807-8

DONATE BLOOD NOW
Schedule an appointment to give blood now!
Go
About our test book

home
about the tour
about blood
being a donor
recipient stories
sponsor a drive
young blood
games & more

SAVE A LIFE TOUR 2003
American Red Cross



## Safety

The Red Cross is committed to protecting the safety of both donors and potential recipients of donated blood.

### Donor Safety

Donating blood is a safe and easy process.

- » Register with Us
  - » Top 10
  - » How To Help
  - » Blood Donation Process
  - » Apheresis
  - » Tips for a Good Donation Experience
  - » Donor Eligibility
  - » Safety
  - » FAQ
  - » Thank You, Donors!
- It is not possible to get AIDS or other infectious disease by giving blood.  
*A sterile, disposable needle is used for each blood donation. Once used, the needle is discarded.*
  - Feeling faint or fatigued after donating blood is rare or minor.  
*If it occurs, it most likely will pass in a matter of hours.*
  - You can only donate if your health history permits and you feel well.  
*You are asked some health questions and are given a mini physical — temperature, pulse, blood pressure, and red cell count check - prior to donation to ensure that you are feeling well and that it is safe for you to give blood.*
  - Your health history and test results are confidential and cannot be given out without your permission, except as required by law.
  - You can help ensure your experience is a positive and rewarding one:  
*Stay in the canteen area for the requested period of time; mention to the staff any unusual feelings or sensations; and avoid strenuous exercise or heavy lifting for about 5 hours after donating.*

### Blood Safety

Assuring the safety of the blood supply is a high-tech process requiring rigorous testing, proper processing, labeling, and storage, and careful quality control. To help ensure that the blood is as safe as possible, the American Red Cross:

- Accepts donations only from voluntary blood donors
- Provides information about high-risk behaviors associated with transmissible diseases that may impact one's ability to donate blood
- Conducts a behavioral and health history interview and a mini physical exam with all donors prior to donation
- Provides a confidential opportunity for donors to ask that their blood be discarded

### How many times have you donated blood?

Submit

Enter a whole number from 0 to 999.

Note: The following information will be based on whole blood donations. It may not be accurate for apheresis donors.





- Provides a confidential 800-number donors can call with any questions or concerns post donation
- Tests every donation\* for infectious diseases, including HIV, hepatitis B and C virus, syphilis, and other infectious diseases, and discards units which have abnormal test results
- Invests in research and technology to support the development of new and more sophisticated tests

#### Tests performed on each unit of donated blood\*

| Disease                                       | Test                             | Year Implemented |
|---|----------------------------------|------------------|
| <b>HIV/ AIDS</b>                              | HIV/AIDS HIV- I Antibody test    | 1985             |
|   | HIV-1/2 Antibody test            | 1992             |
|   | HIV-I p24 Antigen test           | 1996             |
| <b>HIV/ AIDS and Hepatitis C</b>              | Nucleic Acid Test (NAT) **       | 1999             |
| <b>Hepatitis C</b>                            | Hepatitis C Anti-HCV             | 1990             |
| <b>Hepatitis B</b>                            | Hepatitis B Surface Antigen test | 1971             |
|   | Hepatitis B Core Antibody        | 1987             |
| <b>Hepatitis</b>                              | Hepatitis ALT                    | 1986             |
| <b>Syphilis</b>                               | Syphilis Serologic test          | 1948             |
| <b>Human T-cell Lymphotropic Virus (HTLV)</b> | HTLV-I Antibody                  | 1989             |
|   | HTLV -I/II Antibody              | 1998             |


\* Not all tests are performed on autologous donations (blood a recipient donates for him or herself). If an autologous donation is not used by the donor, it is discarded.

\*\*A West Nile Virus test will be added to this list in 2003 under an investigational license.

NOTE: This list is subject to change as new blood safety opportunities and requirements emerge. Additional tests may be performed to meet special patient needs.

© Copyright The American Red Cross. All Rights Reserved.

[RedCross.org](http://www.redcross.org) | [Red Cross Store](http://www.redcross.org) | [Privacy Policy](http://www.redcross.org) | [Blood Information](http://www.redcross.org) | [Contact Us](http://www.redcross.org) | [Search](http://www.redcross.org) | [Links](http://www.redcross.org)



ISSN: 09/616,283; Art Unit: 1645  
Attorney Docket No. VRXB-P01-001

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Application of:

GOODNOW

Serial No: 09/616,283

Filed: July 14, 2000

For: SYSTEM FOR DETECTING  
BACTERIA IN BLOOD, BLOOD  
PRODUCTS, AND FLUIDS OF  
TISSUES

Art Unit: 1645

Attorney Docket No. VRXB-P01-001

Examiner: J. Hines

Assistant Commissioner of Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

**Declaration Under 37 C.F.R. §1.132**

Sir:

I, Harvey G. Klein, MD., hereby declare as follows:

1. I am the Chief of the Department of Transfusion Medicine at the Warren C. Magnuson Clinical Center, National Institutes of Health. I have been working and conducting research in the field of transfusion medicine for over thirty (30) years. I serve on the Food and Drug Administration's (FDA) Blood Products Advisory Committee. I actively hold leadership positions in the American Association of Blood Banks, the American Blood Commission, the American Society of Hematology, and the American Society of Apheresis. I currently serve as Chairman of the Committee of Revision for Blood and Blood Products for the U.S. Pharmacopeia. I am a diplomat of the American Board of Internal Medicine and a diplomat of the American Board of Pathology. I have authored and co-

authored more than 175 publications on transfusion medicine. Accordingly, my curriculum vitae is attached as Appendix A.

2. I have read the above-identified application, the pending claims, and the Office Action mailed on September 29, 2003. I have been asked by Verax Biomedical, Inc. to comment upon the meaning of the term "donor" to a person of skill in this area.
3. In the blood banking discipline, the term "donor" is a universally recognized term of art. Federal Law mandates that only healthy and asymptomatic subjects are allowed to donate blood. Prior to donating, each individual must undergo a thorough screening process that includes temperature, pulse, blood pressure, hemoglobin content in the blood, and questions about general health, background and travel. See AABB Association Bulletin #99-10, dated December 2, 1999, attached herewith as Appendix B. See also the blood donation eligibility guidelines as set forth by the American Red Cross, herewith attached as Appendix C. If the donor fails any of these questions/requirements, the individual is turned away from the donation center to prevent provision of adulterated components to blood recipients. Thus, in the blood banking arena the term "donor" means a healthy or asymptomatic subject qualified to donate blood.
4. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both,



under Section 1001 of Title XVIII of the United States Code and that willful false statements may jeopardize the validity of this Application for Patent or any patent issuing thereon.

Dated: 12/1/2003

Signature: Harvey G. Klein  
Dr. Harvey G. Klein  
Chief, Department of Transfusion Medicine  
Warren C. Magnuson Clinical Center  
National Institutes of Health

## **CURRICULUM VITAE**

**HARVEY G. KLEIN, M.D.**

**DATE OF BIRTH:** May 8, 1943

**PLACE OF BIRTH:** Boston, Massachusetts

**MARITAL STATUS:** Married, June 20, 1965

**WIFE'S NAME:** Sigrid Petri

**CHILDREN:** Susanna Rebecca, born July 26, 1969  
Stephan Eliot, born February 3, 1972  
Melissa Kathryn, born June 30, 1976

**HOME ADDRESS:** 13628 Canal Vista Court, Potomac, MD 20854

**OFFICE ADDRESS:** National Institutes of Health  
Warren G. Magnuson Clinical Center  
Department of Transfusion Medicine  
10 Center Drive, Building 10, Room 1C711  
Bethesda, MD 20892-1184

### **EDUCATION:**

|           |  |
|-----------|--|
| 1955-1961 | Boston Latin School (Franklin Medalist)    |
| 1961-1965 | Harvard College, A.B. (magna cum laude)    |
| 1965-1969 | The Johns Hopkins School of Medicine, M.D. |

### **POSTGRADUATE TRAINING:**

|           |   |
|-----------|---|
| 1969-1970 | Internship, The Johns Hopkins Hospital<br>Department of Medicine            |
| 1970-1972 | Resident, The Johns Hopkins Hospital,<br>Department of Medicine             |
| 1972-1973 | Fellow in Hematology, The Johns Hopkins<br>Hospital, Department of Medicine |

### **MILITARY SERVICE:**

|           |   |
|-----------|---|
| 1973-1975 | Commissioned Corps, Public Health Service |
|-----------|---|

**PROFESSIONAL APPOINTMENTS:**

|              |  |
|--------------|--|
| Present:     | Chief, Department of Transfusion Medicine  |
| 1997-present | Executive Committee, Warren G. Magnuson Clinical Center                                |
| 1990-94      | Special Assistant to the Director for Science, Warren G. Magnuson Clinical Center, NIH |
| 1983         | Tenured Scientist  |
| 1976-83      | Assistant Chief, Blood Bank Department, NIH  |
| 1976-83      | Chief, Blood Services Section, Clinical Center Blood Bank, NIH                         |
| 1975-6       | Acting Chief of Blood Services, Clinical Center Blood Bank, NIH                        |
| 1974-5       | Chief, Hemorrhagic Diseases Program, Division of Blood Diseases, NHLBI, NIH            |
| 1973-5       | Assistant to the Director, Division of Blood Diseases and Resources, NHLBI, NIH        |

**CERTIFICATION:**

Maryland Medical License, 1969 (#K8711)  
Diplomate, American Board of Internal Medicine, 1972 (#39283)  
Diplomate, Hematology Subspecialty - ABIM, 1976 (#39283)  
Diplomate, American Board of Pathology, Blood Bank Specialty, 1977

**ACADEMIC APPOINTMENT:**

1995-present Johns Hopkins School of Medicine – Visiting Professor, Pathology  
1990-present Johns Hopkins School of Medicine – Visiting Professor, Medicine

**PROFESSIONAL SOCIETIES:**

American Federation of Medical Research  
American Society of Hematology  
American Association of Blood Banks  
American Society for Apheresis  
International Society of Blood Transfusion

**AWARDS AND HONORS:**

Detur Award (Harvard)  
Phi Beta Kappa  
Award of Distinction, American Society of Medical Writers, 1975  
Commendation Medal, Public Health Service, 1980

Peacock Lecturer, University of Texas (Dallas), 1984  
 Equal Opportunity Achievement Award, Public Health Service, 1985  
 Elmer L. DeGowin Lectureship, University of Iowa, 1986  
 Meritorious Service Medal, Public Health Service, 1989  
 PHS Unit Commendation, 1991  
 Clarence Schein Memorial Lectureship, Montefiore Medicine Center,  
 New York, 1991  
 Visiting Professor, Mayo Clinic Section of Transfusion Medicine, 1991  
 NIH Clinical Center Director's Award, 1992  
 John B. Alsever Award (contributions to the blood banking sciences), 1993  
 Annual Lecture Award - American Society for Apheresis and 5th World  
 Apheresis Association Congress, 1994  
 Charles E. Walter Memorial Award, Mid Atlantic Association of Blood Banks,  
 1995  
 Allen Latham Award (European Society for Haemapheresis), 1995  
 Tibor J. Greenwalt Lectureship, 1996  
 Cohn DeLaval Medal (World Apheresis Association), 1996  
 Joseph R. Bove Visiting Professorship, Yale, 1997  
 Rita and Taft Schreiber Memorial Lectureship Cedars-Sinai Medical Center, 1998  
 Corresponding Membership, German Blood Transfusion Society, 1998  
 Distinguished Visiting Professor of Pathology, The Johns Hopkins School of  
 Medicine, 1999  
 Visiting Professor of Pathology, University of Pennsylvania, 1999  
 Chairman, 24<sup>th</sup> International Symposium of Blood Transfusion, The  
 Netherlands, 1999  
 HHS Secretary's Distinguished Services Award, 2000  
 NIH Director's Award, 2000  
 Millennial Medical Festival Lecturer, Royal College of Physicians, London, 2000  
 Distinguished Lecturer, Texas Blood Institute, San Antonio, 2000  
 Emily Cooley Award (American Association of Blood Banks) 2001  
 Francis Morrison Memorial Lecture (American Society for Apheresis) 2003  
 Keynote Speaker, Paul Ehrlich Institute (Frankfurt) 2003

#### **CONSULTANT APPOINTMENT:**

Attending Physician, Clinical Hematology Branch, NHLBI, 1975-96  
 FAES Graduate School at NIH (Immunohematology), 1976-1986  
 National Heart, Lung and Blood Institute, Division of Extramural Affairs,  
 1977-present  
 National Blood Resources Education Program (NBREP) 1988-91  
 Uniformed Services University of Health Science, Clinical Asst. Prof.,  
 1977-present  
 Letterman Army Institute of Research, Blood Peer Review Panel, 1987-92  
 Erythropoietin Advisory Board - Ortho Biotech Corp., 1989-95  
 Haemonetics Corp. Scientific Advisory Board

AIBS - Medical Free-electron Laser Program, 1990-91  
 Somatogen Corp., Advisory Board (recombinant human hemoglobin), 1989-98  
 American Medical Association - Drug Evaluations Annual, 1992  
 Cryopharm - Scientific Advisory Board (lyophilized blood cells), 1994-5  
 Zymquest - Scientific Advisory Board (universal red cells), 1994-present  
 AMA Therapeutic Technology Assessment  
 Office of Naval Research – Navy-Army Blood Program, 1998-2000  
 British Blood Authority, 1998 - present  
 Alliance Pharmaceuticals (perfluorocarbon oxygen carriers), 1998-Present  
 New Jersey Department of Health & Senior Services (Bloodless Surgery) 1999  
 Sangart (hemoglobin-based oxygen carriers), 1999-Present  
 World Health Organization, 2000  
 Viacell Medical Scientific Advisory Board (cell expansion) 2000 – Present  
 Vitex Scientific Advisory Board (pathogen reduction of blood) 2001-Present  
 Gambro BCT Medical Advisory Board (pathogen reduction of blood) 2001  
 Johns Hopkins Hospital Advanced Transfusion Practices Center, 2000 – 2002

#### **EDITORIAL BOARD MEMBERSHIP:**

Editor-in-Chief, Journal of Clinical Apheresis, 1986-94  
 Transfusion, 1994-present  
 Transfusion Medicine Reviews, 1994-present  
 Blood, 1988 - 1993  
 Infusion Therapy and Transfusion Medicine (German Society of Blood Transf.)  
 NIH Alumni Update (Founding Member), 1989 - 1996

#### **PROFESSIONAL ACTIVITIES:**

##### **American Association of Blood Banks**

President, 2000-2001  
 President – elect, 1999-2000  
 Vice President, 1997-1999  
 Board of Directors, 1995-2002  
Chairman, Board of Trustees, Research Foundation, 1985-9  
Co-chairman, Scientific Strategic Plan, 1989  
 Transfusion Practices Committee, 1982-8  
 Therapeutic Hemapheresis Committee, 1983-6  
 Nominations Committee, 1990, 1992, 2003  
 Ad Hoc Standards Committee for Marrow Processing Facilities, 1991  
 Strategic Planning Committee, 1991-3,  
 Standards Committee, 1991-6, Chairman, 1993-6  
 North America Task Force for Stem Cell Standards, 1994-6  
 Board of Directors, National Blood Data Resource Center, 2002-3

## **American Blood Commission**

Board of Directors, 1985-8, ASFA representative

## **American National Red Cross**

Blood Services Advisory Committee, 1981-3  
Committee for Protection of Human Subjects, 1983-7  
Scientific Council for Biomedical Research , 1985-9  
Chairman, Transfusion Science Initial Review Group, 1985-9  
Board of Directors, Washington Region , 1989-94  
Chairman, Medical Advisory Committee, Blood Services Washington  
Region, 1988-92  
Blood Services Committee, Washington Region, 1988-90  
Blood Services Committee, Greater Chesapeake and Potomac Region,  
1990  
Board of Directors, Chesapeake and Potomac Blood Program, 1992 -  
present

## **American Society of Hematology**

Scientific Subcommittee in Clinical Laboratory  
Hematology, 1991-97  
Transfusion Medicine Subcommittee, 1993-8  
Chairman, Transfusion Medicine Education Program, 1997, 1998  
International Outreach Initiative – Subcommittee for Asia

## **American Society for Apheresis**

President, 1985  
Board of Directors, 1981-7  
Chairman, Clinical Applications Committee, 1983-7  
Chairman, Scientific Program, 1984, 1990

## **FDA**

Blood Products Advisory Committee, 2002 - present  
Advisory Panel on Orphan Drugs  
Blood donors at risk for AIDS - subject matter expert  
Workshop on Criteria for Evaluation of Red Cell Substitutes, 1999  
Workshop on Leukoreduction of Blood Components, 1999

## **Haemonetics Corporation**

Medical Advisory Board, 1988-99

Board of Directors, 1998-2002

**Institute of Medicine**

Blood Safety Forum, 1994-95  
Committee on Resuscitation Fluids, 1998

**International Society of Blood Transfusion**

Scientific Committee – Int'l Congress (2000) Vienna  
BEST Working Party, 1999-present  
Scientific Committee International Congress (2004 Edinburgh)

**Metropolitan Washington Blood Banks, Inc.**

Board of Directors, 1979-82  
Chairman, Committee on Regionalization, 1979-80

**National Marrow Donor Program (NMDP)**

President, NMDP Council, 1993-4  
Board of Directors, 1992-95  
Membership Committee, 1988-92  
Research and Publications Committee, 1990-95  
International Affairs Committee, 1990-92  
Vice President for Donor Centers, NMDP Council, 1992-94

**National Heart, Lung and Blood Institute**

National Blood Resources Education Program  
Chairman, Data Safety and Monitoring Committee,  
TRAP Study, 1991-6  
Chairman, Data Safety & Monitoring Committee, T-Cell  
Depletion Trial, 1994-2002

**National Institute on Aging**

Clinical Research Subpanel, 1982-5

**National Institute of Allergy and Infectious Diseases**

Expert Panel on the Effects of Donor Pool Size on the Safety and Efficacy  
of Immunoglobulin Products, 1998

**U.S. Public Health Service**

Advisory Committee on Blood Safety and Availability, 2001 - present  
PHS Executive Task Force on AIDS, Charlottesville, VA, 1988  
Medical Review Board, 1982-96  
USPHS Working Group on HTLV-1, 1988  
Medical Board, Clinical Center, NIH, 1982-4, 1989-91  
NIH Strategic Planning Committee (Molecular Medicine  
Initiative Subcommittee), 1992  
Secretary's Committee on Clinical Center Options, 1995

### **U.S. Pharmacopeia**

Committee of Revision, Chairman: Blood and Blood  
Products, 1995-2000  
Committee of Experts – Chairman, Blood & Blood Product Standards,  
2000-2005

### **World Apheresis Association**

Board of Directors, 1993-5  
Councillor - North America, 1986-88, 1992-4  
Scientific Program Committee, 1984, 1986, 1992, 1994

### **World Health Organization**

Consultant to Global Blood Program - 2001

Invited lecturer (including plenary or keynote lectures) at numerous national and international meetings.

Ad hoc reviewer for numerous scientific and medical journals.



## **BIBLIOGRAPHY:**

### **PEER REVIEWED JOURNALS:**

Klein HG, Bell W. Disseminated Intravascular Coagulation Occurring During Heparin Therapy. *Ann Int Med* 1974; 80:477-481.

Klein HG, Aledort LM, Bouma BN., et al. A Cooperative Study for the Detection of the Carrier State of Classic Hemophilia. *N Engl J Med* 1977; 296:959-962.

Klein HG. A Chimpanzee Breeding Colony for Hepatitis Research: *J Med Primat* 1978; 7:182-184.

Weiss GB, Nienhuis AW, McIntosh CL, Klein HG. Traumatic Cardiac Hemolytic Anemia as a Late complication of a Starr-Edwards Mital Valve Prosthesis. *Arch Intern Med* 1979; 139:373-375.

Klein HG, Faltz LA, McIntosh CL, Appelbaum F, Deisseroth AB, Holland PV. Surgical Hypothermia in a Patient with a Cold Agglutinin: Management by Plasma Exchange. *Transfusion* 1980; 20:354-357.

Miller DM, Winslow RM, Klein HG, Wilson KC, Statham NJ, Brown FL, Fulmer J. Improved Exercise Performance after Exchange Transfusion in Subjects with Sickle Cell Anemia. *Blood* 1980;56:1127-32.

Klein HG, Garner RJ, Miller DM, Rosen S, Statham NJ, Winslow RM. Automated Partial Exchange Transfusion in Sickle Cell Anemia. *Transfusion*, 1980; 20:578-84.

Sazama K, Klein HG, Davey RJ, Corash L. Intraoperative Hemolysis as the Initial Manifestation of Glucose-6-Phosphate Dehydrogenase Deficiency. *Arch Intern Med* 1980; 140:845-6.

Kolins J, Allgood JW, Burghardt D, Klein HG, McGinniss MH. Modification of B, I, i, and Lewis Antigens in a patient with DiGuglielmo's erythroleukemia. *Transfusion* 1980; 20:574-77.

Diamond WJ, Brown F, Bitterman P, Klein HG, Davey R, Winslow RM. Delayed Hemolytic Transfusion Reaction Presenting as Sickle Cell Crisis. *Ann Int Med* 1980; 93:231-34.

- Coles SM, Klein HG, Holland PV. Alloimmunization in two multi-transfused patient populations: Sick Cell Disease and Thalassemia Major. *Transfusion* 1981; 21:462-67.
- Corash L, Klein HG, Deisseroth A, Shafer B, Rosen S, Beaman J, Griffith P, Nienhuis A. Selective Isolation of Young Erythrocytes for Transfusion Support of Thalassemia Major Patients. *Blood* 1981; 57:599-606.
- Gadek JE, Klein HG, Holland PV, Crystal RG. Replacement Therapy of Alpha 1-antitrypsin Deficiency: Reversal of Protease-Imbalance within the Alveolar Structures of Piz Subjects. *J Clin Invest* 1981; 68:1158-65.
- Kessler CV, Klein HG, Havlik RJ. Uncontrolled Thrombocythemia in Chronic Myeloproliferative Disorders. *Brit J Haem* 1982; 50(1):157-67.
- Klein HG, Kessler C, Anderson KC, Loomis KF. Multiple Arterial and Venous Thromboemboli in a Male with Chronic Myelogenous Leukemia and Occult Metastatic Breast Carcinoma. *South Med J*. 1982; 75:745-747.
- Lenes B, Klein HG, Lakatos E. Selective Removal of Sick Cells with the IBM 2997 Continuous Flow Blood Cell Separator. *J. Clin. Apheresis* 1983; 1:64-70.
- Young NS, Klein HG, Griffith P, Nienhuis AW. A trial of immunotherapy in aplastic anemia and pure red cell aplasia. *J. Clin. Apheresis* 1983; 1:95-103.
- Reichert CM, Weisenthal LM, Klein HG. Delayed Hemorrhage Following Percutaneous Liver Biopsy. *J. Clin. Gastroenterol.* 1983; 5:263-266.
- Bracey A, Klein HG, Chambers S, Corash L. Simplified Selective Isolation of Young Red Cells for Chronic Transfusion Support. *Blood* 1983; 61:1068-1071.
- Winslow RM, Butler WM, Kark JA, Klein HG, Moo-Penn W. The effects of bloodletting on exercise performance in a subject with a high-affinity hemoglobin variant. *Blood* 1983; 62: 1159-1164.
- Klein HG, McCurdy PR, Braine HG, MacPherson JL, Dowling RN. Centrifugal Leukocytapheresis and Thrombocytapheresis: Complications Related to Equipment and Technique. *J. Clin. Apheresis* 1983; 1:190-195.
- Rose J, Klein H, Greenstein J, McFarlin D, Gerber L, McFarland H. Lymphocytapheresis in chronic progressive multiple sclerosis: Results of a preliminary trial. *Ann Neurol* 1983; 14:593-594.
- Marti GE, Folks T, Longo DL, Klein HG. Concise review: Therapeutic Cytapheresis in chronic lymphocytic leukemia. *J Clin Apheresis* 1983; 1:243-248.

Dowling R, Weber V, Osborne L, Klein HG. Mononuclear cell collection using various techniques. *J Clin Apheresis* 1984; 2:32-40.

Lane HC, Masur H, Longo DL, Klein HG, Rook AH, Quinnan GV, Steis RG, Macher A, Whalen G, Edgar LC, Fauci AS. Partial Immune Reconstitution in a Patient with the Acquired Immunodeficiency Syndrome. *N Engl J Med* 1984; 311: 1099-1109.

Rodgers GP, Bonner RF, Klein HG, Noguchi CT, Nienhuis AW, Schechter AN. Periodic Microcirculatory Flow in Patients with Sickle Cell Disease. *N Engl J Med* 1984; 311: 1534-8.

Grindon AJ, Tomasulo PA, Bergin JJ, Klein HG, Miller JD, Mintz PD. The Hospital Transfusion Committee: Guidelines for Improving Services. *JAMA* 1985; 253:540-543.

Winslow RM, Klein HG, Monge CC. Red cell mass in Andean natives with excessive polycythemia. *Arch Biol Andina* 13:85-94, 1984-85.

Winslow RM, Monge CC, Brown EG, Klein HG, Sarnquist F, Winslow NJ. The effect of hemodilution on O<sub>2</sub> transport in high altitude polycythemia. *J Appl Physiol* 1985; 59(5): 1495-1502.

Tomasulo PA, Lenes BA, Noto TA, Klein HG, Menitove J. Automatic Special Case Consultations in Transfusion Medicine. *Transfusion*, 1986; 26:186-193.

Klein HG, Balow JE, Dau PC, Hamburger MI, Leitman SF, Pineda AA, and Tindal RSA: Clinical Application of Therapeutic Apheresis. *J Clin Apheresis* 1986; 3(7):1-93.

AuBuchon JP, Carter CS, Adde MA, Meyer DR, Klein HG. Optimization of Parameters for Maximization of Plateletpheresis and Lymphocytophoresis yields on the Haemonetics Model V50. *J. Clin. Apheresis* 1986; 3(2):103-110.

Stevenson HC, Stevenson GW, Leitman SF, Carter CS, Alvord G, Klein HG. The isolation of human mononuclear subsets by cytapheresis for laboratory research. *Plasma Ther Transfus Technol* 1986; 7:365-371.

Muul LM, Nason-Burchenal K, Carter CS, Cullis H, Slavin D, Hyatt C, Director EP, Leitman SF, Klein HG, Rosenberg SA. Development of an automated closed system for generation of human lymphokine activated killer (LAK) cells for use in adoptive immunotherapy. *J Immunol Methods* 1987; 101:171-181.

Carter CS, Leitman SF, Cullis H, Muul LM, Nason - Burchenal K, Rosenberg SA, Klein HG. Use of a continuous flow cell separator in density gradient isolation of lymphocytes. *Transfusion* 1987; 27:362-365.

- Areman EM, Simonis T, Carter C, Read EJ, Klein HG. Bulk cryopreservation of lymphocytes in glycerol. *Transfusion* 1988; 28:151-156.
- Rodgers GP, Schechter AN, Noguchi CT, Klein HG, Nienhuis AW, Bonner RF. Microcirculatory adaptations in sickle cell anemia. Reactive hyperemia response. *Am J. Physiol* 1990;258:
- Kruskall MS, Mintz PD, Bergin JJ, Johnston MFM, Klein HG, Miller JD, Rutman R, Silberstein L. Transfusion therapy in emergency medicine. *Ann Emergency Med* 1988; 17:327-335.
- Sloand EM, Kessler CM, McIntosh CL, Klein HG. Neutralization of heparin-induced anticoagulation with methylene blue. *Thromb. Res.* 1989, 54:677-86.
- Carter CS, Leitman SF, Cullis H, Muul LM, Nason-Burchenal K, Rosenberg SA, Klein HG. Technical Aspects of lymphokine-activated killer cell production. *J. Clin. Apheresis* 1988; 2/3:113-117.
- Leitman SF, Klein HG, Melpolder JJ, Shih JW, Read EJ, Harvarth L, Darr F, Foy JL, Alter HJ. Clinical implications of positive tests for antibodies to human immunodeficiency virus type 1 in asymptomatic blood donors. *N Engl J Med* 1989; 321:917-23.
- Sloand EM, Fox SM, Banks SM, Klein HG. Preparation of IgA deficient platelets. *Transfusion* 1990; 30:322-326.
- Loiacono B, Carter G, Leitman S, Klein HG. Efficacy of various methods of confidential unit exclusion in identifying potentially infectious blood donations. *Transfusion* 1989; 29:823-6.
- DePalma L, Palmer R, Leitman SF, Dolan WD, Klein HG. Utilization patterns of frozen autologous red blood cells: experience in a referral center and a community hospital. *Arch Pathol Lab Med* 1990; 114:516-18.
- Bergsten P, Amitai G, Kehrl J, Dhariwal KR, Klein HG, Levine M. Millimolar concentrations of ascorbic acid in purified human mononuclear leukocytes: depletion and reaccumulation. *J Biol Chem* 1990; 265:2584-2587.
- Sloand EM, Klein HG. Effect of white cells on platelets during storage. *Transfusion* 1990; 30:333-338.
- Lane HC, Zurich KM, Wilson W, Cefali F, Easter M, Kovacs JA, Masur H, Leitman SF, Klein HG, Steis RG, Longo DL, Fauci AS. Syngeneic bone marrow transplantation and adoptive transfer of peripheral blood lymphocytes combined with zidovudine in human immunodeficiency virus infection. *Ann Int Med* 1990; 113:512-19.

Alter HJ, Epstein JS, Swenson SG, Van Raden MA, Klein HG, Ward JW, Kaslow RA, Menitove JE, Sandler SG, Sayers MH, Hewlett IK, Chernoff AI. Prevalence of Human Immunodeficiency Virus Type 1 (HIV-1) p24 antigen in U.S. Blood Donor: Assessment of Efficacy for Blood Donor Screening. *N Engl J Med* 1990; 323:1312-17.

Sloand EM, Sloand JA, Prodouz K, Klein HG, Yu MW, Harvath L, Fricke W. Reduction of platelet glycoprotein Ib in uremia. *Brit J. Haem* 1991; 77:375-81.

Fricke W, Alling D, Kimball J, Griffith P, Klein H. Lack of efficacy of tranexamic acid in thrombocytopenic bleeding. *Transfusion* 1991; 31:345-348.

Keilholz U, Klein HG, Korbling M, Brado B, Carter CS, Cullis H, Galin F, Hunstein W. Peripheral blood mononuclear cells collection from patients undergoing adoptive immunotherapy or peripheral blood-derived stem cell transplantation and from normal donors. *J. Clin Apheresis* 1991; 6:131-6.

Sloand EM, Klein HG, Pastakia KP, Pierce P, Prodouz KN. Effects of albumin on the inhibition of platelet aggregation by beta lactam antibiotics. *Blood* 1992; 79:2022-27.

Sloand EM, Klein HG, Banks SM, Vareldzis B, Merritt S, Pierce P. Epidemiology of thrombocytopenia in HIV infection. *Eur J. Haem* 1992; 48:168-172.

Natanson C, Hoffman WD, Koev LA, Dolan DP, Banks SM, Bacher J, Danner RL, Klein HG, Parrillo JE. Plasma exchange does not improve survival in a canine model of human septic shock. *Transfusion* 1993; 33:243-8.

Stroncek DF, Holland PV, Bartsch G, Bixby T, Simmons RG, Antin JH, Anderson KC, ASH PC, Bolwell BH, Hansen JH, Heal JM, Henslee-Downey PJ, Jaffe' ER, Klein HG, Lau PM, Perkins HA, Popovsky MA, Price TH, Rowley SD, Stehling LC, Weiden PL, Wissel ME, McCullough J. Experiences of the first 493 unrelated marrow donors in the National Marrow Donor Program. *Blood* 1993; 81:1940-46.

Sloand E, Kumar P, Klein HG, Merritt S, Sacher R. Transfusion of blood products in HIV-1 infected persons: relationship to opportunistic infection. *Transfusion* 1994; 34:48-53.

Sloand EM, Kumar P, Yu M, Klein HG. Effect of amphotericin B and fluconazole on platelet membrane glycoproteins. *Transfusion* 1994; 34: 415-20.

Sloand EM, Alyono D, Klein HG, Chang P, Yu M, Lightfoot FG, Kessler C. 1-Deamino-8-D-arginine vasopressin (DDAVP) increases platelet membrane expression of glycoprotein 1b in patients with disorders of platelet function and after cardiopulmonary bypass. *Am J. Hematol* 1994; 46: 199-207.

Lubarsky DA, Hahn C, Bennett DH, Smith LR, Bredehoeft SJ, Klein HG, Reves DG. The hospital costs (fiscal year 1991/1992) of a simple perioperative allogeneic red blood cell transfusion at Duke University. *Anesth Analg* 1994; 79:629-37.

Lee JH, Klein HG. The effect of donor erythrocyte sedimentation rate on efficiency of granulocyte collection by centrifugal leukapheresis. *Transfusion* 1995; 35:384-388.

Lee JH, Cullis H, Leitman SF, Klein HG. The efficacy of pentastarch in granulocyte collection by centrifugal leukapheresis. *J Clin Apheresis*. 1995;10:198-202.

Lee JH, Leitman SF, Klein HG. A controlled comparison of the efficacy of hetastarch and pentastarch in granulocyte collections of centrifugal leukapheresis. *Blood* 1995; 86:4662-6.

Sloand EM, Alyono D, Yu M, Klein HG. Platelet membrane glycoproteins and microvesicles in blood from post-operative salvage: a study in cardiac bypass patients. *Transfusion* 1995; 35:738-744.

Blaese RM, Culver KW, Miller AD, Carter CS, Fleisher T, Clerici M, Shearer G, Chang L, Chiang Y, Tolstosher P, Greenblatt J, Rosenberg S, Klein HG, Berger M, Mullen C, Ramsey J, Muul L, Morgan RA, Anderson WF. T lymphocyte-directed gene therapy for ADA Deficiency (SCID): Initial trial results after 4 years. 1995; *Science* 270:475.

Sloand EM, Yu M, Klein HG. Comparison of random-donor platelets prepared from whole blood units and single-donor platelet's prepared from single donor apheresis collections. *Transfusion* 1996; 36:955.

Read EJ, Kunitake ST, Carter CS, Chau Q, Klein HG. Enumeration of CD34+ hematopoietic progenitor cells in peripheral blood and leukapheresis products by microvolume fluorimetry: a comparison with flow cytometry. *J Hematotherapy*. 1997, 6:291-301.

Walker RE, Carter CS, Muul L, Natasajan V, Herpin BR, Leitman SF, Klein HG, Mullen CA, Baseler M, Falloon J, Davey Jr RT, Kovacs JA, Polis MA, Masur H, Blaese RM, Lane HC. Peripheral expansion of mature T Cells is a major means of CD4+ T cell regeneration in adults with HIV-1 infection. *Nature Med*. 1998, 4:852-6.

Silva N, O'Bryan L, Medeiros E, Holand H, Suliman J, Patronas N, Reed SG, Klein HG, Masur H, Badaro R, Trypanosoma Cruzi meningoencephalitis in HIV-infected patients. *J. Acquir Immun Defic Syndr Hum Retrovirol* 1999; 20:342-9.

Zumberg MS, Procter JL, Lottenberg R, Kitchens CS, Klein HG. Autoantibody formation in the alloimmunized red blood cell recipient. *Arch Int Med* 2001; 161:281-90.

Stroncek DF, Procter JL, Moses L, Bolan C, Pomper GJ, Conry-Cantilena C, Malech HL, Klein, HG, Leitman, SF. Intravenous Rh immune globin prevents alloimmunization in D negative granulocyte recipients, but obscures the detection of an alloanti-K. *Immunology* 2001, 17:37-41.

Wang E, Adams S, Panelli M, Klein H, Marincola FM. Fluorimetric detection of single nucleotide variants (SNV) by proportional hybridization to oligonucleotide arrays (PHOA). *Nature Biotechnology* (in press)

Stroncek DF, Rainer T, Sharon V, et al. Sick cell hemoglobin polymerization in red cell components from donors with sick cell trait prevents effective leukocyte reduction by filtration. *Transfusion* 2002;42:1466-72

Panelli MC, Wang E, Phan G, Puhlman M, Miller L, Ohnmacht GA, Klein HG, Marincola FM. Gene expression profiling of the response of peripheral blood mononuclear cells and melanoma metastases to systemic IL-2 administration. *Genome Biol* 2002; 25:3(7) Research 0035

Haddad SA, Winer KK, Gupta A, Chakrabarti S, Noel P, Klein HG. A puzzling case of anemia. *Transfusion* 2002;42:1610-13.

Wang E, Adams S, Zhao Y, Panelli MC, Simon R, Klein H and Marincola FM. A strategy for detection of known and unknown SNP using a minimum number of oligonucleotides applicable in the clinical settings *Journal of Translational Medicine* 2003, 1:4 (online; 20 Aug 2003)

Leitman SF, Browning JM, Yau YY, Mason G, Klein HG, Conry-Cantilena C, Bolan CD. Impact of using hemochromatosis subjects as allogeneic blood donors: a prospective study. *Transfusion* (in press)

Guralnick J, Woodman R, Klein HG et al. Anemia and functional status in an elderly population. *JAMA* (in press)

## **REVIEWS:**

Borberg H, Dahlke MB, Fraser ID, Fratantoni JC, French JE, Holland PV, Klein HG, Polesky HF, Sandler SG. Which are the established or potential risks for Donors undergoing Cytopheresis Procedures and how can they be prevented? *Vox Sang* 1980; 39(3):169-79.

Klein HG. Selective Isolation of Young Red Cells for Chronic Transfusion. *Plasma Therapy* 1981; 2:175-80.

Klein HG, Winslow RW. Partial Exchange Transfusion in Sick Cell Disease. *Prog Clin Biol Res* 1982; 88:235-243.

Klein HG. Transfusions with young erythrocytes (neocytes) in Sick Cell Anemia. *Am J Ped Hemat Onc* 1982; 4(2):162-165.

Klein HG. Therapeutic Cytapheresis. *Prog Clin Biol Res* 1982; 106:297-306.

Klein HG. Therapeutic Apheresis: An overview. In: Nose, Y., Malchesky, P.S., and Smith, J.W. (Eds). *Plasmapheresis* ISAO Press (Cleveland) 1983:3-6.

Klein HG. Therapeutic Cytapheresis. In: Kolins J, Jones J, eds. *Therapeutic Apheresis*, Washington: American Association of Blood Banks, 1983:23-34.

Klein HG. The Kinship of Coagulation and Immunity: The Contact System. In Tregellas WM and Keating LJ, eds. *Immunology*. Arlington, VA: American Association of Blood Banks, 1985:177-190.

Klein HG, Alter HJ. Blood Transfusion and AIDS In: *AIDS*, Chicago: American Medical Association 1987:7-10.

Sloand EM and Klein HG. Therapeutic Apheresis. In: Kasprisin DO (ed) *Current Status of Hemapheresis*, Washington: American Association of Blood Banks 1987:9-47.

Klein HG, Leitman SF. Adoptive immunotherapy in malignant disease. *Transfusion* 1989; 29:170-178.

Higgins MJ, Klein HG. Massive transfusion in the intensive care unit. *J Intensive Care Med* 1989; 4:221-233.

The use of autologous blood. The National Blood Resource Education Program Expert Panel. *JAMA* 1990; 263:414-17.

Klein HG. Controversies in transfusion medicine: Alanine aminotransferase screening of blood donors: Pro. *Transfusion* 1990; 30:363-367.

Klein HG. When is Transfusion the best option? *Emergency Medicine*. 1992; 24(5): 59-66.

Strauss RG, Ciavarella D, Gilcher RO, Kasprisin DO, Kiproff DD, Klein HG, McLeod BC. Clinical Application of Therapeutic Hemapheresis. *J. Clin Apheresis* 1993; 8:189-269.

Walker R, Blaese RM, Carter CS, Chang L, Klein H, Lane HC, Leitman SF, Mullen CA, Larson M. A study of the safety and survival of the adoptive transfer of genetically marked syngeneic lymphocytes in HIV-1 infected identical twins. *Human Gene Ther* 4 (5) 1993; 659-80.



Klein HG. Immunologic Aspects of Blood Transfusion. Seminars in Oncology 1994; 21:16-20.

Lee JH, Klein H.G. Mononuclear cell adoptive immunotherapy Hematology/Oncology Clinics of North America 1994; 8(6):1203-1221.

Lee JH, Klein HG. Peripheral blood stem cell transplantation Hematology/Oncology Clinics of North America 1994; 9(1):1-22.

Lee JH, Klein HG. Cellular gene therapy. Hematology/Oncology Clinics of North America 1994; 9(1):91-113.

Sloand EM, Pitt E, Klein HG. Safety of the blood supply. JAMA 1995; 274:1368-73.

Klein HG. Allogeneic transfusion risks in surgical patients. Am J. Surg 1995; 170 (suppl): 215-265.

Hillyer CD, Klein HG. Immunotherapy and gene transfer in the treatment of the oncology patient: Role of transfusion medicine. Transf Med Rev 1996; 10(1):1-8.

Miller Y, Klein HG. Growth factors and their impact on Transfusion Medicine. Vox Sang 1996; 71:196-204.

Klein HG. New insights into the management of anemia in the surgical patient. Am J Med 1996; 101 (2A): 125-155.

Klein HG, Strauss RG, Schiffer CA. Granulocyte transfusion therapy. Semin Hematol 1996; 33:359-368.

Klein HG, Dodd RY, Dzik WH, Luban NLC, Ness PM, Pisciotto P, Schiff P, Snyder EL. Current status of solvent/detergent-treated frozen plasma. Transfusion 1998; 38:102-107.

Klein HG. Immunologic Aspects of Transfusion: A once and future risk? Anesthesiology 1999; 91:861-5.

Engelfriet CP, Reesink HW, Blajchman MA et al. Bacterial contamination of blood components. Vox Sanguinis 2000; 78:59-67.

Engelfriet CP, Reesink HW, Klein HG, et.al. International Forum: granulocyte concentrates. Vox Sang 2000; 79:59-66.

Klein HG. The immunomodulatory effects of blood transfusion. Tumori 2001; 87(2) 18-20

Engelfriet CP, Reesink HW, Klein H.G. et al. The future of pathogen-in concentrates. Vox Sang 2003; 85:54-66.

Brittenham GM, Klein HG, Kushner JP, Ajioka RS. Preserving the nation's blood supply. Hematology (Am Soc Hematol Educ Program). 2001; 422-32.

#### **BOOK CHAPTERS:**

Klein HG. Effects of plasma exchange on plasma constituents: Choice of solutions. In: MacPherson JL and Kasprisin DO (Eds). Therapeutic Plasma Exchange. Plenum Press Inc. (Boca Raton) Vol II, 1985:3-14.

Klein HG. Erythrocytapheresis. In: Valbonesi M, Pineda AA, Biggs JC (Eds). Hemapheresis, Milan: Wichtig Editore. 1986:143-148.

Read EJ, Klein HG. Hematological Effects of Aging: Considerations for the Elderly. In: Cutler NR, Narang PK, eds. Drug Studies in the Elderly: Methodology and Applications. New York: Plenum 1986:123-150.

Klein HG. Blood Product Therapy in the Critical Care Unit. Parrillo JE (Ed). Therapy in Critical Care Medicine, Philadelphia: B.C. Decker Inc. 1988:143-148.

Klein HG. Transfusion support of hemoglobinopathies. In: Cash J (ed). Transfusion Medicine Vol III. Churchill Livingstone (Edinburgh) 1987:143-148.

Pineda AA, Klein HG. Applications of Hemapheresis. In: Greenwalt T (Ed). Hematology, Vol 17. Blood Transfusion, New York: Churchill Livingstone (Edinburgh).1988: 213-233.

Klein HG. Young red cells for transfusion support. In: Pineda AA, Valbonesi M, eds. Donor Hemapheresis, Milan, Italy: Wichtig Editore. 1986:143-148.

Lacerna LV, Hester J, Pineda AA, Burgstaler E, Klein HG. Clinical/Transfusion support in adoptive cellular immunotherapy: The role of cytapheresis, In: Stetler-O'Gara K (Ed) Adoptive Cellular Immunotherapy of Cancer, New York: Marcel Dekker 1990: 175-190.

Klein HG, Travis LB. Transfusion support of hemoglobinopathies. In: Modern Transfusion Therapy, Vol I, CRC Press Inc., Boca Raton, 1990: 143-148.

Klein HG. Apheresis. In: Hoffman R, Benz EJ, Shattil S, Furie B, Coates T (Eds). Hematology: Principles and Practice. New York: Churchill Livingstone 1990: p1638-43.

Klein HG. Transfusions with young erythrocytes (neocytes) in Sick Cell Anemia. *Am J Ped Hemat Onc* 1982; 4(2):162-165.

Klein HG. Therapeutic Cytapheresis. *Prog Clin Biol Res* 1982; 106:297-306.

Klein HG. Therapeutic Apheresis: An overview. In: Nose, Y., Malchesky, P.S., and Smith, J.W. (Eds). *Plasmapheresis* ISAO Press (Cleveland) 1983:3-6.

Klein HG. Therapeutic Cytapheresis. In: Kolins J, Jones J, eds. *Therapeutic Apheresis*, Washington: American Association of Blood Banks, 1983:23-34.

Klein HG. The Kinship of Coagulation and Immunity: The Contact System. In Tregellas WM and Keating LJ, eds. *Immunology*. Arlington, VA: American Association of Blood Banks, 1985:177-190.

Klein HG, Alter HJ. Blood Transfusion and AIDS In: *AIDS*, Chicago: American Medical Association 1987:7-10.

Sloand EM and Klein HG. Therapeutic Apheresis. In: Kasprisin DO (ed) *Current Status of Hemapheresis*, Washington: American Association of Blood Banks 1987:9-47.

Klein HG, Leitman SF. Adoptive immunotherapy in malignant disease. *Transfusion* 1989; 29:170-178.

Higgins MJ, Klein HG. Massive transfusion in the intensive care unit. *J Intensive Care Med* 1989; 4:221-233.

The use of autologous blood. The National Blood Resource Education Program Expert Panel. *JAMA* 1990; 263:414-17.

Klein HG. Controversies in transfusion medicine: Alanine aminotransferase screening of blood donors: Pro. *Transfusion* 1990; 30:363-367.

Klein HG. When is Transfusion the best option? *Emergency Medicine*. 1992; 24(5): 59-66.

Strauss RG, Ciavarella D, Gilcher RO, Kasprisin DO, Kiproff DD, Klein HG, McLeod BC. Clinical Application of Therapeutic Hemapheresis. *J. Clin Apheresis* 1993; 8:189-269.

Walker R, Blaese RM, Carter CS, Chang L, Klein H, Lane HC, Leitman SF, Mullen CA, Larson M. A study of the safety and survival of the adoptive transfer of genetically marked syngeneic lymphocytes in HIV-1 infected identical twins. *Human Gene Ther* 4 (5) 1993; 659-80.

Klein HG. Immunologic Aspects of Blood Transfusion. Seminars in Oncology 1994; 21:16-20.

Lee JH, Klein H.G. Mononuclear cell adoptive immunotherapy Hematology/Oncology Clinics of North America 1994; 8(6):1203-1221.

Lee JH, Klein HG. Peripheral blood stem cell transplantation Hematology/Oncology Clinics of North America 1994; 9(1):1-22.

Lee JH, Klein HG. Cellular gene therapy. Hematology/Oncology Clinics of North America 1994; 9(1):91-113.

Sloand EM, Pitt E, Klein HG. Safety of the blood supply. JAMA 1995; 274:1368-73.

Klein HG. Allogeneic transfusion risks in surgical patients. Am J. Surg 1995; 170 (suppl): 215-265.

Hillyer CD, Klein HG. Immunotherapy and gene transfer in the treatment of the oncology patient: Role of transfusion medicine. Transf Med Rev 1996; 10(1):1-8.

Miller Y, Klein HG. Growth factors and their impact on Transfusion Medicine. Vox Sang 1996; 71:196-204.

Klein HG. New insights into the management of anemia in the surgical patient. Am J Med 1996; 101 (2A): 125-155.

Klein HG, Strauss RG, Schiffer CA. Granulocyte transfusion therapy. Semin Hematol 1996; 33:359-368.

Klein HG, Dodd RY, Dzik WH, Luban NLC, Ness PM, Pisciotto P, Schiff P, Snyder EL. Current status of solvent/detergent-treated frozen plasma. Transfusion 1998; 38:102-107.

Klein HG. Immunologic Aspects of Transfusion: A once and future risk? Anesthesiology 1999; 91:861-5.

Engelfriet CP, Reesink HW, Blajchman MA et al. Bacterial contamination of blood components. Vox Sanguinis 2000; 78:59-67.

Engelfriet CP, Reesink HW, Klein HG, et.al. International Forum: granulocyte concentrates. Vox Sang 2000; 79:59-66.

Klein HG. The immunomodulatory effects of blood transfusion. Tumori 2001; 87(2) 18-20

Engelfriet CP, Reesink HW, Klein H.G. et al. The future of pathogen-inactivated platelet concentrates. Vox Sang 2003; 85:54-66.

Brittenham GM, Klein HG, Kushner JP, Ajioka RS. Preserving the national blood supply. Hematology (Am Soc Hematol Educ Program). 2001; 422-32

### **BOOK CHAPTERS:**

Klein HG. Effects of plasma exchange on plasma constituents: Choice of replacement solutions. In: MacPherson JL and Kasprisin DO (Eds). Therapeutic Hemapheresis CRC Press Inc. (Boca Raton) Vol II, 1985:3-14.

Klein HG. Erythrocytapheresis. In: Valbonesi M, Pineda AA, Biggs JC, eds. Therapeutic Hemapheresis, Milan: Wichtig Editore. 1986:143-148.

Read EJ, Klein HG. Hematological Effects of Aging: Considerations for Clinical Trials. Cutler NR, Narang PK, eds. Drug Studies in the Elderly: Methodological Concerns, New York: Plenum 1986:123-150.

Klein HG. Blood Product Therapy in the Critical Care Unit. Parrillo JE, ed. Current Therapy in Critical Care Medicine, Philadelphia: B.C. Decker Inc. 1987:281-287.

Klein HG. Transfusion support of hemoglobinopathies. In: Cash J (ed) Progress in Transfusion Medicine Vol III. Churchill Livingstone (Edinburgh) 1987: 198-213.

Pineda AA, Klein HG. Applications of Hemapheresis. In: Greenwalt TJ (Ed). Methods in Hematology, Vol 17. Blood Transfusion, New York: Churchill Livingstone (Edinburgh). 1988: 213-233.

Klein HG. Young red cells for transfusion support. In: Pineda AA, Valbonesi M, Sniecinski IJ, eds. Donor Hemapheresis, Milan, Italy: Wichtig Editore, 1988: p. 65-69.

Lacerna LV, Hester J, Pineda AA, Burgstaler E, Klein HG. Clinical/Technical challenges in adoptive cellular immunotherapy: The role of cytappheresis, In: Stevenson HC (Ed) Adoptive Cellular Immunotherapy of Cancer, New York: Marcel Dekker, Inc. 1989: 175-190.

Klein HG, Travis LB. Transfusion support of hemoglobinopathies. In: Dutcher JP (ed) Modern Transfusion Therapy, Vol I, CRC Press Inc., Boca Raton, 1990: p. 93-106.

Klein HG. Apheresis. In: Hoffman R, Benz EJ, Shattil S, Furie B, Cohen HJ, eds. Hematology: Principles and Practice. New York: Churchill Livingstone Inc. 1990: p1638-43.

Klein HG, Apheresis in transition. In: Nydegger UE, (ed): Therapeutic Hemapheresis in the 1990's. Curr Stud Hematol Blood Transf. Basel, Karger, 1990, No. 57, pp 1-9.

Klein HG. Plasma exchange in the future: innovations and new indications. In Rossi EC, Simon TL and Moss GS (eds): Principles of Transfusion Medicine. Williams and Wilkins, Baltimore 1991: p557-560.

Klein HG. Therapeutic mononuclear cell transfusion: adoptive immunotherapy. In Rossi EC, Simon TL, and Moss GS (eds): Principles of Transfusion Medicine, Williams and Wilkins, Baltimore. 1991: p295-300.

Klein HG. Autologous mononuclear cell therapy. In Taswell HF and Pineda AA (eds): Autologous Transfusion and Hemotherapy. Blackwell Scientific Publication, Boston. 1991: p220-232.

Leonard K, Klein HG. Acute Hemolytic Anemia. In Bell WR (ed): Hematologic and Oncologic Emergencies. Contemporary Management in Critical Care. Churchill Livingstone, New York 1993: pp33-69.

Klein, H.G. Immunohematology and Blood Transfusion. In Isselbacher KJ, Braunwald E, Wilson JD, Martin JB, Fauci AS, Kasper DL (eds): Harrison's Principles of Internal Medicine, McGraw-Hill Inc, New York 1993: p 2377.

Klein HG. Principles of Apheresis. In Anderson KC and Ness PM (eds). Scientific Basis of Transfusion Medicine. WB Saunders, Philadelphia 1994: p 781-796.

Cantilena CC, Klein HG. Therapeutic use of blood components. In Rakel CT (ed): Conn's Current Therapy. W.B. Saunders Company, Philadelphia, PA 1994: p 427-31.

Higgins MJ, Klein HG. Blood Products in the ICU. In Parrillo JE and Bone RC (eds): Critical Care Medicine: Principles of Diagnosis and Management. Mosby-Year Book, Inc. Philadelphia, PA 1995, p 1323-44.

Klein HG. Hemapheresis and Cellular Therapy. In Hoffman R, Benz EJ, Shattil S, Furie B, Cohen HJ (eds): Hematology: Principles and Practice, New York: Churchill Livingstone Inc. 1995: p2038-2045.

Klein HG. Therapeutic Hemapheresis. In: Petz LD, Swisher S, Kleinman S, Spence R, Strauss RG, (eds): Clinical Practices of Transfusion Medicine. Churchill Livingstone, New York 1996, p1011-1022.

Carter CS, Gress R, Klein HG. Routine methods for preparing hematopoietic progenitor cell components. In Lasky L and Warkentin P (eds): Marrow and Skin Cell Processing for Transplantation. American Association of Blood Banks, Bethesda, MD 1995, p 51-68.

Klein HG. Immunohematology, In: Rich RR, Fleisher TA, Schwartz BD, Shearer WT, Strober W (eds): Clinical Immunology: Principles and Practice. Mosby, St. Louis 1995, p2220-2228.

Klein HG. Novel Cellular Therapies, In: Van Oss CJ (ed). Transfusion Immunology and Medicine. Marcel Dekker, Inc. (New York) 1995; 411.

Klein HG. The immunomodulatory effects of blood transfusion. In: Petz LD, Swisher S, Kleinman S, Spence R, Strauss RG (eds): Clinical Practice of Transfusion Medicine. Churchill Livingstone, New York 1996, p 59-70.

Klein HG. Therapeutic Hemapheresis. In: Petz LD, Swisher S, Kleinman S, Spence R, Strauss RG, (eds) : Clinical Practices of Transfusion Medicine. Churchill Livingstone, New York 1996, p 1011-1022.

Lee JH, Klein HG. Therapeutic mononuclear cell transfusion: adoptive immunotherapy and gene therapy. In: Rosse EC, Simon TL, and Moss GS (eds): Principles of Transfusion Medicine. Williams and Wilkins, Baltimore 1996, p 329-44.

Higgins MS, Klein HG. Blood Products in the Critical Care Unit. Parrillo JE ed. Current Therapy in Critical Care Medicine. Mosby-Year Books, Inc. Philadelphia 1997, p 362-7.

Klein HG. Cellular Gene Therapy. In McLeod B, Price TH, Drew MJ (eds): Principles of Apheresis. AABB Press, 1997, p 537-46 .

Klein HG. Continuing issues regarding transfusion and coagulation factor-acquired HIV infection. In: Pizzo PA, Wilfert CM (eds); Pediatric AIDS. (3<sup>rd</sup> Edition) Baltimore, Williams and Wilkins, 1998.

Lee JH, Klein HG. Mononuclear cell transfusion: Immunotherapy Using Allogeneic Donor-Derived Lymphocytes. Mintz PD (ed) Transfusion Therapy: Clinical Principles and Practice. AABB Press, Bethesda 1998, p 251-265.

Klein HG. Principles of Apheresis. In: Anderson KC and Ness PM (eds) Scientific Basis of Transfusion Medicine. WB Saunders, Philadelphia 1999: p553-68.

Conry-Cantilena, Klein, HG. Hemochromatosis subjects as blood and tissue donors. In: Barton JC and Edwards CO, (eds); Hemochromatosis. Cambridge University Press, Cambridge 2000: p564-73.

Lazarus EF, Klein HG. Hemapheresis and cellular therapy. In: Hoffman R. Benz EJ, Shattil S, Furie B, Cohen HJ, Silberstein LE, McGlave P (eds): Hematology: Principles and Practice. Churchill Livingstone Inc.: New York 1999, p2291-99.

Klein HG. Molecular Biology in Transfusion Medicine. In: Smit Sibinga C Th and Klein HG (eds). Molecular Biology in Transfusion Medicine. Kluwer Academic Publishers (Dordrecht) 2000, p3-12..

Kicklighter EJ, Klein HG. Hemolytic transfusion reactions. In: Linden JV, Bianco (eds). Blood Safety and Surveillance, Marcel Dekker, inc., New York 2001, p47-70..

Lee, J-H, Klein HG. From leukocyte reduction to leukocyte transfusion: the immunologic effects of transfused leukocytes. Contreras M (ed) Bailliere Tindal, London 2000, p585-600.

Griffith L, Klein HG. Immunohematology, In: Rich RR, Fleisher TA, Kotzin B, Shearer WT, Schroeder H (eds): Clinical Immunology: Principles and Practice. Churchill Livingstone, London 2001, p127.1-127.10.

Lazarus EF, Klein HG. Apheresis, In Rich RR, Fleisher TA, Kotzin B, Shearer WT, Schroeder H (eds): Clinical Immunology: Principles and Practices. Churchill Livingstone, London, 2001, p118.1-118.9

Bolan CD, Klein HG. Transfusion Medicine and pharmacologic aspects of hemostasis. In Kitchens CS, Alving BM, Kessler CM (eds): Consultative Hemostasis and Thrombosis. W.B. Saunders Company, New York 2002, 395-418.

Klein HG. The evolution of cellular therapy in the new millennium. In: Sacher RA (ed) Cellular Therapy: New Frontiers in transfusion Medicine. AABB Pres (Bethesda) 2002, 1-13.

Klein HG. Cellular gene therapy. In: McLeod BC (ed) Apheresis: Principles and Practice. AABB Press (Bethesda) In Press

Lazarus EF, Klein HG. Hemapheresis and cellular therapy. In: Hoffman R, Benz EJ, Shattil S, Furie B, Cohen HJ, Silberstein LE, McGlave P (eds): Hematology: Principles and Practice. Churchill Livingstone Inc.: New York (in press)

## **GOVERNMENT PUBLICATIONS:**

Petit CR, Klein HG. Hemophilia, Hemophiliacs and the Health Care System. DHEW Publ. No. (NIH) 1975:76-781.

Klein HG. Isovolemic Hemodilution in High Altitude Polycythemia. In: Adjustment to High Altitude, DHHS Publ. No. (NIH) 1983; 83-2496: 47-52.



Anderson D, Epstein J, Pierik L, Solomon J, Blattner W, Saxinger L, Alter H, Klein H, McCurdy P, Nemo G, Kaplan J, Allen J, Khabbaz R, Lairmore M. Licensure of screening tests for antibody against human T-lymphocytotropic virus type I. MMWR 37(48):736-740, December 9, 1988.

Toy PT, Dzik W, Gould SA, Hoppe PA, Kruskall MA, Maurer RC, Menitove JE, Naiman JL, Thurer RL, Klein HG, McCurdy PR, Sloand EM. Use of Autologous Blood. U.S. Department of Health and Human Services Publ. No. (NIH) 89-3038.

Stehling LC, Cosgrove DM, Moss GS, Schiffer CA, Slichter SJ, Snyder EJ, Klein HG, McCurdy PR, Sloand EM. Indications for the use of red blood cells, platelets, and fresh frozen plasma. U.S. Department of Health and Human Services. Publ. No. (NIH) 89-2974.

Toy PT, Beattie C, Gould SA, et al. Transfusion Alert. Use of Autologous Blood. U.S. Department of Health and Human Services Publ. No. (NIH) 94-3038.

#### **PUBLISHED PROCEEDINGS:**

Klein HG, Alter HJ, Holland PV. Postexposure immunoglobulin prophylaxis of hepatitis B: A comparison of two dosage schedules. In Vyas GN, Cohen SM, Schmid R (Ed): Viral Hepatitis: Proceedings of the Second Symposium of Viral Hepatitis (Franklin Institute Press, Philadelphia) 1978:483-86.

Klein HG. Neocytapheresis: Collection, Preparation and Preliminary Clinical Results, in Proceedings of the Red Cross Apheresis Symposium, Chicago, American Red Cross Blood Services, Mid America Region. 1979:88-96.

Klein HG. Red Cell Exchange Transfusion in Secondary Polycythemia. In Proceedings of the Red Cross Second Apheresis Symposium, Chicago, 1980:74-82.

Spiro T, Young N, Klein HG. Collection of Erythroid (BFU-e) and Myeloid (CFU-c) Progenitor Cells from Peripheral Blood using Continuous Flow and Intermittent Flow Cell Separators. in Proceedings of the Red Cross Second Apheresis Symposium, Chicago, 1980; 158-72.

Klein HG. Cell Separators for Red Cell Exchange in Advances in the Pathophysiology, Diagnosis, and Treatment of Sickle Cell Disease (Alan R. Liss, Inc., New York) 1982; 109-117.

Schulz SC, van Kammen DP, Waters R, Klein HG, Balow JE, Bunney WE. Double Blind Evaluation of Plasmapheresis in Schizophrenic Patients: A Pilot Study. Artificial Organs 1983; 12:3141.

- Young NS, Klein HG, Griffith P, Nienhuis AW. Therapeutic plasma exchange and lymphocyte depletion in aplastic anemia and pure red cell aplasia. In: Nose, Y., Malchesky, P.S., J.W., and Krakauer, R.S. (Eds). Plasmapheresis Raven Press (New York) 1983:339-348.
- Funck-Brentano JL, Klein HG, Lysaght MJ, Oda T. Plasma Exchange or Plasma Treatment. In: Nose Y., Malchesky A.S. and Smith J.W. (eds). Plasmapheresis: New Trends in Therapeutic Applications. ISAO Press (Cleveland) 1983: 167-171.
- Groopman JE, Klein HG, Levine PH, Sande MA, Redfield RR. AIDS: Questions and Controversies - A Panel Discussion. Sem Hem 1987; 24 (3) Suppl 2: 9-19.
- Klein HG. Adoptive immunotherapy: novel applications of blood cell separators. J. Clin Apheresis 1988; 4:198-202.
- Chernoff AI, Klein HG, Sherman LA (Eds). Research opportunities in transfusion medicine. Transfusion 1989; 29:711-742.
- Stevenson HC, Klein HG. Biologic response modifiers (BRM) chemotherapy and activated cytotoxic leukocyte subsets: strategies for rational design of apheresis-based cancer immunotherapies. Prog Clin Biol Res 1990; 337:559-65.
- Immunosuppressive effect of blood transfusion. Klein HG and Tarrter PI (eds) Dept of Surgery, Mt. Sinai Medical Center (New York) 1991.
- Klein HG. Current risks of blood transfusion. In Friedel N, Hetzer R, Royston D, (eds) Blood Use in Cardiac Surgery. Steinkopff Verlag (Darmstadt) 1991:69-73.
- Klein HG. Future Technologies in Transfusion Medicine. Infusionstherapie 1991; 18 (suppl 1):31-34.
- Shih JW-K, Kung K, Dragon EA, Davies C, Pottathil R, Leitman SF, Klein H, Spadaro J, Melpolder J, Alter HJ. Quantitative analysis of an HIV positive blood donor cohort. In Andrieu JM (ed) Viral Quantitation in HIV Infection. John Libbey Eurotext, (Paris) 1991; 171-176.
- Klein HG. Transfusion Medicine: The impact of biotechnology, growth factors, and bioengineering. In Smit Sibinga C. Th, Das PC, and Cash JD. Transfusion Medicine: Fact and Fiction. Kluwer Academic Publishers (Dordrecht) 1991;153-158.
- Cottler-Fox M, Read EJ, Yu M, Carter C, Klein HG. Mononuclear and CD34 stem cell recovery after automated ficoll processing of marrow or peripheral blood for transplantation. In: Gross S, Gee AP, Worthington-White D eds. Prog Clin Biol Res, New York: Wiley-Liss, 1992; 377:569-73.

Klein HG. Transfusion in Transplant Patients: the Good, the Bad and the Ugly. *Journal of Heart/Lung Transplantation* 1993; 12(S):7-12.

Klein HG. Apheresis, immunoreconstitution and AIDS. In Agishi T (ed) *Therapeutic Plasmapheresis (XII)* VSP Publishers (Tokyo) 1993; 21-23.

Cottler-Fox M, Klein HG. Transfusion support of hematology and oncology patients. The role of hematopoietic growth factors. *Arch Pathol Lab Med* 1984; 118:417-20.

Klein HG. Oxygen Carriers and transfusion medicine. *Art Cells Blood Subs and Immob Biotech* 1994; 22:123-135.

Klein HG. Novel cellular therapies. *Immunol Invest* 1995; 24:411-20.

Klein HG. Cellular Gene Therapy: An overview. *J. Clin Apheresis* 1994; 9:139-41.

Klein HG. Immunologic aspects of blood transfusion. *Semin Oncol* 1994; 21 (2), Suppl 3, 16-20.

Klein HG, McCurdy PR, Nemo G, et al. Transfusion alert: Use of autologous blood. *Transfusion* 1995; 35:703-11.

Klein HG. Examples of risks that we tolerate. In: Bendixen H, Manning FJ, Sparacino L (eds). *Blood and Blood Products: Safety and Risk*. National Academy Press (Washington), 1996, p101-105.

Klein HG, Dodd RY, Ness PM, Fratantoni JA, Nemo GJ. Current status of microbial contamination of blood components: summary of a conference. *Transfusion* 1997; 37:95-101.

#### **BOOKS EDITED:**

*Standards for Blood Banks and Transfusion Services*, 16th Edition, American Association of Blood Banks, Bethesda 1994.

*Standards for Blood Banks and Transfusion Services*, 17th Edition, American Association of Blood Banks, Bethesda, MD, 1996.

*Transfusion Medicine*. *Semin Hematol* (Guest Editor), 1996; 33 (4) : 275-358.

*Molecular Biology in Transfusion Medicine*. Smit Sibinga C Th and Klein HG (eds). Kluwer Academic Publishers (Dordrecht) 2000.

#### **EDITORIALS:**

- Klein HG. Therapeutic Plasma Exchange: A Healthy Dose of Skepticism. *Western J Med* 1983;92-94.
- Klein HG. Blood Transfusion and Athletics; Games People Play. *New Engl J Med* 1985; 312:854-856.
- Klein HG. Transfusion Medicine: The evolution of a new discipline. *JAMA* 1987; 258: 2108-2109.
- Klein HG. Developmental Milestones: The Journal at five years. *J Clin Apheresis* 1988, 4:1-2.
- Klein HG. Wolf in Wolf's clothing: Is it time to raise the bounty on the passenger lymphocyte? *Blood* 1992; 80:1865-8.
- Klein HG. It seemed a shame to throw away the red cells. Selective component collection. *Transfusion* 1993; 33:788-90
- Klein HG. Preface. In: Manning FG and Sparacino L (eds): *Blood Donors and the Supply of Blood and Blood Products*. National Academy Press (Washington) 1996, p ix-x.
- Klein HG. Introduction: Transfusion Medicine. *Semin Hematol* 1996; 33: 275-6.
- Blajchman MA, Klein HG. Looking back in anger: Retrospection in the face of a paradigm shift. *Transf Med Rev*. 1997 ; 11 (1) :1-5.
- Klein HG. Cellular Gene Therapy: A Credible Odyssey. *Transfusion*. 1997; 37:675-77.
- Klein HG. Blood, Drugs, and the United States Pharmacopeia. *Transf Med Rev* 1998; 12(2):81-3.
- Klein HG. Transfusion Safety: Avoiding unnecessary bloodshed. *Proc. Mayo Clin* 2000, 75: 5-7.
- Klein HG. The prospect of red-cell substitutes. *N Engl J Med* 2000; 342: 1666-8.
- Klein HG. Will blood transfusion ever be safe enough? *JAMA* 2000; 284(2): 238-240.
- Klein HG. Blood donation: It's more than comfort and convenience. *AABB News* 2001; January/February: 2.
- Klein HG. The transfusion timeline and current good health care practices. *AABB News* 2001; May/June: 2

Klein HG. Safety and availability: The CJD paradigm. AABB News 2001; July/August:2.

Klein HG. Rushing into the new millennium. AABB News 2001; September/October: 1

Klein HG. Earthquake in American. Transfusion 2001. 41:1179-80.

Klein HG. Older is not necessarily better. Anesthesiology 2003 98(4):807-8

**ASSOCIATION BULLETIN**  
**#99-10**

**Date:** December 2, 1999

**To:** AABB Institutional Members

**From:** Paul Ness, MD                      Karen Shoos Lipton, JD  
              President                              Chief Executive Officer

**Re:** New Uniform Donor History Questionnaire Issued

The uniform donor history questionnaire has been prepared by the AABB Blood Bank/Transfusion Service Standards Program Unit of the Standards Program Committee. The AABB Board of Directors has charged this committee to update the questionnaire as needed to be consistent with AABB Standards and Food and Drug Administration (FDA) requirements. This version of the questionnaire supersedes the existing questionnaire, which was issued on June 12, 1998, as Association Bulletin #98-3. It was amended on February 16, 1999, in Association Bulletin #99-5. This revision complies with the 19th edition of *Standards for Blood Banks and Transfusion Services*.

The donor screening process is an important tool that is designed to help safeguard the nation's blood supply. It is imperative that screening be performed consistently to prevent unsuitable donor candidates from donating. The process must be comprehensive, comprehensible and educational.

The questions have been prepared to assist in developing uniform guidelines for blood donor screening nationwide. They have been carefully worded to convey content simply and briefly, and have been reorganized into logical groupings. Changes made since the last version reflect the deliberations and decisions of the Standards Program Committee and recent statements issued by the FDA. New questions have been added, cites have been updated and comments have been revised so it is recommended that blood establishments review this questionnaire in its entirety.

The FDA has reviewed and approved the questionnaire, which is in compliance with the current FDA regulations/recommendations for donor suitability. The FDA further stated that "when distributing this questionnaire to your membership, it would be prudent to remind them to use it *in toto, without modifications*." (Emphasis added by the AABB.)

Licensed blood establishments need to report changes in the donor history questionnaire in accordance with 21 CFR 601.12 (Changes to an approved application). Implementation of the FDA-approved AABB uniform donor history questionnaire without modification is reported in an establishment's annual report (21 CFR 601.12(d)).

You may be aware that the donor screening process has come under increasing public scrutiny, and has been discussed at recent meetings of the FDA Blood Products Advisory Committee and the Public Health Service's Advisory Committee on Blood Safety and Availability. These discussions have focused on the need for uniform donor screening throughout the country.

Although the questionnaire is not an AABB requirement and is provided as a tool to help members with compliance on donor screening, the AABB strongly recommends that each member institution administer the questions, as written, as part of its donor screening process.

\*AABB members may also view this document by going to the AABB Web site: [www.aabb.org](http://www.aabb.org). In the Member section of the Web page, click on Library and see the Association Bulletin section.

\*\* The first Association Bulletin of each calendar year provides a listing of Association Bulletins. For copies of AABB association policy statements, including the Technical Bulletins and Joint Statements, fax a request to the Executive Secretary in the AABB Executive Office, (301) 907-6895.

---

| Donor History Questions  | American Association of Blood Banks (AABB)  | Food and Drug Administration (FDA)   | Comments  |
|--|---|--|---|
| 1. Have you ever donated or attempted to donate blood using a different (or another) name here or anywhere else? | Blood collection facilities shall confirm donor identity and link the donor to existing donor records. (Standard B2.100)  | A record shall be available from which unsuitable donors may be identified so those products from such individuals will not be distributed. (21 CFR 606.160(e) April 1999)   |   |
| 2. In the past 8 weeks, have you given blood, plasma or platelets here or anywhere else?                         | Frequency of whole blood donation is every eight weeks. (Standard B1.300)   | Frequency of blood donation is every 8 weeks unless otherwise approved by the medical director (21 CFR 640.3(f) April 1999)  | Infrequent plasma donors can donate every four weeks. (FDA Memos 3/10/95 <sup>1</sup> and 12/14/95 <sup>2</sup> )                       |
| 3. Have you for any reason been deferred or refused as a blood donor or told not to donate blood?                | No specific requirement.  | No specific requirement.   |   |
| 4. Are you feeling well and healthy today?   | The prospective donor shall appear to be in good health. (Standard B2.000)  | Donor must be determined to be in good general health. (21 CFR 640.3(b) April 1999)  | Requires "yes" answer that tests donor's attention to question context. Initiates sequence of personal health history questions (4-13). |
| 5. In the past 12 months have you been under a doctor's care or had a major illness or surgery?                  | No specific requirements.   | Persons who have received a transfusion of whole blood or a blood component within the past 12 months should not donate blood or blood components. (FDA Memo 4/23/92 <sup>3</sup> )  |   |
| 6. Have you ever had chest pain, heart disease, recent or severe respiratory disease?                            | Prospective donors with diseases of the heart or lungs should be excluded unless determined to be suitable to donate by the blood bank medical director. (Standard B1.700)  | Donor must be free of acute respiratory disease. (21 CFR 640.3(b)(4) April 1999)   |   |
| 7. Have you ever had cancer, a blood disease or a bleeding problem?  | Prospective donors with a history of cancer or abnormal bleeding tendency shall be excluded unless determined to be suitable to donate by the blood bank medical director. (Standard B1.700)  | Persons with hemophilia or related clotting disorders who have received clotting factor concentrates must not donate blood or blood components. (FDA Memo 4/23/92 <sup>3</sup> )   |   |
| 8. Have you ever had yellow jaundice, liver disease, viral hepatitis or a positive test for hepatitis?           | Prospective donors with diseases of the liver shall be excluded unless determined to be suitable to donate by a blood bank physician. (Standards B1.700) Donors with a history of hepatitis after their 11 <sup>th</sup> birthday or a confirmed test for HbsAg or a repeatedly reactive test for anti-HBc are indefinitely deferred. (Standard B2.711) | No individual with a history of hepatitis shall be source of whole blood donation. (21 CFR 640.3(c) April 1999) Exemptions for history of hepatitis before age 11. (FDA Memos 4/23/92 <sup>3</sup> and 12/22/93 <sup>5</sup> ) |   |
| 9. Have you ever had malaria, Chagas' disease or   | Prospective donors who have had a diagnosis of  | Prospective donors who have had malaria should   |   |



|  |  |   |  |
|--|--|---|--|
| babesiosis?  | malaria shall be deferred for 3 years after becoming asymptomatic. (Standard B2. 741) A history of babesiosis or Chagas' disease shall be cause for indefinite deferral. (Standard B2.750)   | be deferred for 3 years after becoming asymptomatic. (FDA Memo 7/26/94 <sup>6</sup> )   |  |
| 10. A. Have you ever taken etretinate (Tegison) for psoriasis?<br><br>B. In the past 3 years, have you taken Acitretin (Soriatane)?<br><br>C. In the past 36 hours, have you taken aspirin, or anything that has aspirin in it?<br><br>D. In the past month, have you taken Isotretinoin (Accutane) or finasteride (Propecia)? | A. People who have received etretinate shall be deferred indefinitely. (Standard B2.520)<br><br>B. Donor is to be Deferred for 3 years From date of last use. (Standard B2. 520)<br><br>C. Ingestion within 36 hours of donation of medications known to irreversibly damage platelet function (eg, aspirin-containing medications) or that inhibit platelet function and have a prolonged half-life should preclude the use of donor as the sole source of platelets for a recipient. (Standard B2.510)<br><br>D. For Accutane, Proscar, or Propecia, donor is to be deferred for 1 month after receipt of last dose. (Standard B2.520) | A. A donor who has taken or is taking Tegison should be permanently deferred. (FDA Memo 7/28/93 <sup>7</sup> )<br><br>B. Donor is to be deferred for 3 years from date of last dose (per manufacturer's insert.)<br><br>C. No specific requirement for whole blood donation. Donors who have recently taken medication containing aspirin, especially within 36 hours, may not be suitable donors for platelet pheresis. (FDA Guidelines 10/7/88 <sup>8</sup> )<br><br>D. A donor taking Accutane or Proscar should be deferred from donating blood for at least one month after receipt of the last dose. (FDA Memo 7/28/93 <sup>7</sup> )<br><br>A donor taking Propecia should be deferred for one month. (FDA Telephone Communication to AABB, January 1998.) | A. Potentially teratogenic: may be present up to 3 years after last use.<br><br>B. Potentially teratogenic. May be present up to 3 years after last use.<br><br>C. Preferred time varies. May be mandated by state health and safety code.<br><br>D. Medication questions grouped. |
| E. In the past 4 weeks, have you taken any pills or medications?<br><br>1. Have you at any time since 1980 injected bovine (beef) insulin?   | E. Drug therapy shall be evaluated by a qualified person to determine suitability to donate blood. (Standard B1.900)   | E. Facility medical director to determine donor acceptability or deferral based on medications. (FDA Memo 7/28/93 <sup>7</sup> )<br><br>1. As a precaution, the FDA recommends that blood donors who have injected bovine insulin since 1980 be indefinitely deferred unless it has been established that the   | E. Medication questions grouped.<br><br>If donor answers "yes" to this question, see #1 below.<br><br>1. If the donor answers "yes" or "don't know" to the question, the FDA recommends that the donor be indefinitely deferred, unless it has been established that the product   |

|   |  |  |  |
|---|--|--|--|
|   |  | product was not manufactured since 1980 from cattle in the United Kingdom. (FDA Guidance for Industry, 11/23/99 <sup>h</sup> )   | was not manufactured since 1980 from cattle in the United Kingdom.<br><br>The FDA recommends that blood establishments review their policies regarding acceptance of insulin dependent diabetic patients as donors and their donor history questions to determine if or at which point in the interview process this question should be asked. The AABB recommends, however, that blood establishments maintain the current order to ensure uniformity of the questionnaire. |
| 11. In the past 4 weeks, have you had any shots or vaccinations?                        | Donors must be queried about vaccines and immunizations. (Standard B2.600)   | No specific requirement.   |  |
| 12. In the past 12 months, have you been given rabies shots?                            | Donor is deferred for 12 months after vaccine for rabies if immunization is given after bite or other exposure to a potentially rabid animal. (Standard B2.600)  | No specific requirement.   |  |
| 13. Female donors: In the past 6 weeks, have you been pregnant or are you pregnant now? | Existing pregnancy or pregnancy in the past 6 weeks is cause for deferral. (Standard B1.800)   | No specific requirement.   |  |
| 14<br>A. In the past 3 years, have you been outside the United States or Canada?        | A. Residents of countries in which malaria is not considered endemic but who have been in an area in which malaria is considered endemic may be accepted as regular blood donors one year after return irrespective of the receipt of antimalarial prophylaxis. (Standard B2.743) Immigrants, refugees or citizens coming from a country in which malaria is considered endemic may be accepted as blood donors 3 years after departure. (Standard B2.742) | A. Travelers to an area considered endemic for malaria should not be accepted as donors of whole blood and blood components prior to one year after departure. After one year, donors free of unexplained symptoms suggestive of malaria may be accepted whether or not they have received antimalarial chemoprophylaxis. Immigrants, refugees and citizens of endemic countries should not be accepted as donors prior to 3 years after departure. After 3 years, donors free of unexplained symptoms suggestive of malaria may be accepted. (FDA Memo 7/26/94 <sup>h</sup> ) | A. Initiates sequence of exposure-type questions (14-30).  |
| B. Have you visited or lived in the United  |  | B. The FDA believes that donors who have   | B. The FDA recommends that within a facility,  |

|  |  |  |   |
|--|--|--|---|
| Kingdom (England, Northern Ireland, Scotland, Wales, the Isle of Man or the Channel Islands) from 1980 to 1996? If so, have you spent a total of six months or more from 1980 through 1996?  |  | resided in the United Kingdom (as identified by these questions) may be at risk for acquiring nvCJD. As a precaution, the FDA recommends that donors who answer "yes" to this question be indefinitely deferred. (FDA Guidance for Industry 11/23/99 <sup>9</sup> )  | current donors need only be questioned once, and new donors questioned at first donation only.  |
| 15<br>A. Have you ever received human pituitary-derived hormone?<br><br>B. Have you received a dura mater (or brain covering) graft?<br><br>C. Have you or any of your blood relatives ever had Creutzfeldt-Jakob disease or have you ever been told that your family is at an increased risk for Creutzfeldt-Jakob disease? | A. Prospective donors who have a family history of Creutzfeldt-Jakob disease or who have received tissue or tissue derivatives known to be a possible source of Creutzfeldt-Jakob agent (eg, dura mater, pituitary growth hormone of human origin) shall be deferred indefinitely. (Standard B2.410) | A. The FDA recommends that any donor who has received injections of pit-hGH be permanently deferred. (FDA Memo 7/28/93 <sup>7</sup> , Guidance for Industry 11/23/99 <sup>9</sup> .)<br><br>B. The FDA recommends that persons who have received transplants of dura mater be permanently deferred from donation. (FDA Guidance for Industry 11/23/99 <sup>9</sup> .)<br><br>C. The FDA recommends that donors at increased risk for CJD be indefinitely deferred and appropriately counseled. The FDA considers that donors who answer "yes" to any of these questions (15 a, b, or c) are at an increased risk for developing CJD. (FDA Guidance for Industry 11/23/99 <sup>9</sup> .) | A. If the donor is uncertain about his or her treatment, the following question may be asked: Was the hormone treatment given repeatedly by injection? Donors who answer "yes" should be deferred.<br><br>C. If the donor is not familiar with the term Creutzfeldt-Jakob disease, this may be taken as a negative response. If the donor is deferred because of family history (one or more family members with CJD), that donor may be re-entered if they meet FDA reentry criteria. (FDA Guidance for Industry 11/23/99 <sup>9</sup> ) |
| 16. In the past 12 months, have you had close contact with a person with yellow jaundice or viral hepatitis, or have you been given Hepatitis B Immune Globulin (HBIG)?  | Close contact with person who has viral hepatitis is a 12 month deferral. (Standard B2.724)  | Close contact with person who has viral hepatitis is a 12 month deferral. (FDA Memo 4/23/92 <sup>10</sup> )  | Close contact generally refers to cohabitation. The medical director should establish a policy for these potential donors. The FDA does not recommend deferral of a sexual partner of an HCV antibody positive individual. (FDA Communication to AABB, August 1999)   |
| 17. In the past 12 months, have you taken (snorted)  | Intranasal use of cocaine is a 12-month deferral.  |  |   |

|   |   |   |                                  |
|---|---|---|----------------------------------|
| cocaine through your nose?  | (Standard B2.727)   |   |                                  |
| 18. In the past 12 months, have you received blood or had an organ or tissue transplant or graft?   | Prospective donors who during the preceding 12 months received blood, blood components or derivatives, or other human tissue known to be possible sources of blood-borne pathogens, shall be excluded. (Standards B2.420, B2.430)   | Persons who have received a transfusion of whole blood or a blood component within the past 12 months should not donate blood or blood components. (FDA Memo 4/23/92 <sup>3</sup> )   | Includes immunization with RBCs. |
| 19. In the past 12 months, have you had a tattoo applied, ear or skin piercing, acupuncture, accidental needlestick or come in contact with someone else's blood?               | Prospective donors shall be deferred from donating blood or blood components for transfusion who, within the preceding 12 months, have a history of: 1) A tattoo. 2) Mucous membrane exposure to blood. 3) Nonsterile skin penetration with instruments or equipment contaminated with blood or body fluids. 4) Sexual or household contact with an individual with viral hepatitis. 5) Sexual contact with an individual with HIV or at high risk of HIV infection. (Standards B2.721, B2.722, B2.723, B2.724, B2.725) | Persons who have had any contact with blood and body fluids through percutaneous inoculation (such as injury or accidental needlestick) or through contact with an open wound, non-intact skin or mucous membrane during the preceding 12 months should be deferred. (FDA Memo 4/23/92 <sup>3</sup> ) |                                  |
| 20.<br>A. In the past 12 months, have you had a positive test for syphilis?<br><br>B. In the past 12 months, have you had or been treated for syphilis or gonorrhea?            | A history of syphilis or gonorrhea, treatment for either, or a reactive screening test for syphilis shall be cause for deferral for 12 months after completion of therapy. (Standard B2.340)  | Persons who have had, or have been treated for, syphilis or gonorrhea during the preceding 12 months should not donate blood or blood components. Persons with a positive (STS) test should be deferred 12 months. (FDA Memo 12/12/91 <sup>11</sup> )   |                                  |
| 21. In the past 12 months, have you given money or drugs to anyone to have sex with you?  | Donor must be given educational material on AIDS high-risk activity, and such at-risk persons should refrain from donating blood. Donor screening shall   | Men and women who have engaged in sex for money or drugs since 1977 and persons who have engaged in sex with such people during the   |                                  |
|   | Include questions intended to identify persons at high risk for HIV infection and high risk for HIV transmission. Such high-risk persons shall be deferred as specified in FDA recommendations. (Standards B3.100, B2.730)  | preceding 12 months should not donate blood or blood components. (FDA Memo 4/23/92 <sup>3</sup> )   |                                  |
| 22.<br>A. At any time since 1977, have you taken money or drugs for sex?<br><br>B. In the past 12 months, have you had sex, even once, with anyone who has taken money or drugs | Refer to question #21   | Men and women who have engaged in sex for money or drugs since 1977 and persons who have engaged in sex with such people during the preceding 12 months should not donate blood or blood components. (FDA Memo 4/23/92 <sup>3</sup> )   |                                  |

|  |   |   |  |
|--|---|---|--|
| for sex?   |   |   |  |
| 23.<br>A. Have you ever used a needle, even once, to take drugs that were not prescribed by a doctor?  | A. Stigmata of narcotic habituation is an indefinite deferral. The donor shall not have used a needle even once to take drugs other than those prescribed by his/her physician. (Standard B2.330) | A. Donor must be free from skin punctures or scars indicative of addiction to self-injected narcotics. (21 CFR 640.3(b)(7) April 1999) Past or present intravenous drug users should not donate blood or blood components. (FDA Memo 4/23/92 <sup>3</sup> ) |  |
| B. In the past 12 months, have you had sex, even once, with anyone who has used a needle to take drugs not prescribed by a doctor?                       | B. Refer to question #21.   | B. Persons who have had sex with any person who is a past or present intravenous drug user should not donate blood or blood components for 12 months. (FDA Memo 4/23/92 <sup>3</sup> )  |  |
| 24. Male donors: Have you had sex with another male, even once, since 1977?  | Refer to question #21.  | Men who have had sex with another man, even one time, since 1977 should not donate blood or blood components permanently. (FDA Memo 4/23/92 <sup>3</sup> )  |  |
| 25. Female donors: In the past 12 months, have you had sex with a male who has had sex, even once, since 1977 with another male?                         | Refer to question #21.  | Females who have had sex with men who have had sex with another man even one time since 1977 should not donate blood or blood components for 12 months. (FDA Memo 4/23/92 <sup>3</sup> )  |  |
| 26.<br>A. Have you ever taken clotting factor concentrates for a bleeding problem, such as hemophilia?   | No specific requirement.  | A. Persons with hemophilia or related clotting disorders who have received clotting factor concentrates should not donate blood or blood components. (FDA Memo 4/23/92 <sup>3</sup> )   |  |
| B. In the past 12 months, have you had sex, even once, with anyone who has taken clotting factor concentrates for a bleeding problem such as hemophilia? |   | B. Persons who have had sex with any person with hemophilia or related clotting disorders who have received clotting factor concentrates should not donate blood or blood components for 12 months. (FDA Memo 4/23/92 <sup>3</sup> )                        |  |
| 27.<br>A. Do you have AIDS or have you had a positive test for the AIDS virus?   | Refer to question #21.  | A. Persons with clinical or laboratory evidence of HIV infection must not donate blood or blood components. (FDA Memo 4/23/92 <sup>3</sup> )  |  |

|   |   |   |   |
|---|---|---|---|
| B. In the past 12 months, have you had sex, even once, with anyone who has AIDS or has had a positive test for the AIDS virus?  |   | B. Persons who have had sex with persons with clinical or laboratory evidence of HIV infection should not donate blood for 12 months. (FDA Memo 4/23/92 <sup>3</sup> )                                  |   |
| 28. Are you giving blood because you want to be tested for HIV or the AIDS virus?   | No specific requirement.                    | No specific requirement.  | Direct question to further evaluate donation motive.  |
| 29. Do you understand that if you have the AIDS virus, you can give it to someone else even though you may feel well and have a negative AIDS test?   | No specific requirement.                    | Donors should be informed that there is an interval during early infection when the HIV antibody test may be negative although the infection may still be transmitted. (FDA Memo 4/23/92 <sup>3</sup> ) | Queries donor understanding of "Important Information for Donors."  |
| 30.<br>A. Were you born in, have you lived in, or have you traveled to any African country since 1977?<br><br>C. When you traveled to <country(ies)> did you receive a blood transfusion or any other medical treatment with a product made from blood?<br><br>D. Have you had sexual contact with anyone who was born in or lived in any African country since 1977? | (Association Bulletin #97-5 <sup>13</sup> ) | (FDA Memo 12/11/96 <sup>12</sup> )  | A. If "no," proceed to the question (question c) about sexual contact.<br><br>If "yes," the donor should be asked to name the specific country(ies). If the donor identifies an African country <i>not</i> listed in the FDA Memo, proceed to question C. If one or more of the countries listed in the FDA Memo is named by the donor, determine if the donor was born in, lived in, or traveled to the country(ies) named by the donor. If the donor was born in or lived in any of the FDA-identified countries, defer indefinitely; questioning stops here. If travel was the donor's risk, ask question B.<br><br>The Central African Republic was the Central African Empire in the late 1970s.<br><br>None of the other countries listed in the FDA Memo have undergone a change in name since 1977.<br><br>Blood establishments should critically evaluate the potential donor's history and statements, and decide whether the individual could have been in the country long enough to have encountered those local conditions related to risk, such as use of unsterile needles or sexual contact.<br><br>When donors report demographic HIV-1 Group O risks, no follow-up actions |
| 30. Continued   |   |   |   |

|   |  |  |   |
|---|--|--|---|
|   |  |  | <p>regarding previously donated blood are necessary.</p> <p>A. If "no," proceed to Question c.</p> <p>If "yes," defer indefinitely.</p> <p>B. If "no," or the donor names a country not identified in the FDA Memo, no deferral.</p> <p>If "yes," ask the donor to specify which country(ies). If donor names country listed in the FDA Memo, defer indefinitely.</p> |
| 31. In the past 12 months, have you been in jail or prison?   | Donors are deferred for 12 months if in the preceding 12 months they have been incarcerated in a correctional institution (jail or prison) for more than 72 consecutive hours. (Standard B2.726) | Individuals who have been incarcerated for more than 72 consecutive hours during the previous 12 months should be deferred as donors for 12 months from the last date of incarceration. (FDA Memo 6/8/95 <sup>14</sup> )   |   |
| 32. Have you read and understood all the donor information presented to you, and have all your questions been answered? | No specific requirement.   | Information should be written in language that assures that the donor understands the definition of high-risk behavior and the importance of self-exclusion. Donors should not be considered suitable unless information about risks can be communicated in the language appropriate to each donor and is constructed to be culturally sensitive to promote comprehension. (FDA Memo 4/23/92 <sup>15</sup> ) |   |

Standards referred to are from the 19<sup>th</sup> edition of *Standards for Blood Banks and Transfusion Services*, effective June 1, 1999.

1. FDA Memorandum, March 10, 1995: Revision of FDA Memorandum of August 27, 1982: Requirements for Infrequent Plasma Donors.
2. FDA Memorandum, December 14, 1995: Donor Deferral Due to Red Blood Cell Loss During Collection of Source Plasma.
3. FDA Memorandum, April 23, 1992: Revised Recommendations for the Prevention of HIV Transmission by Blood and Blood Products.
4. FDA Memorandum, April 23, 1992: Exemptions to Permit Persons with a History of Viral Hepatitis Before the Age of Eleven Years to Serve as Donors of Whole Blood and Plasma: Alternative Procedures, 21 CFR 640.120.
5. FDA Memorandum, December 22, 1993: Donor Suitability Related to Laboratory Testing for Viral Hepatitis and a History of Viral Hepatitis.
6. FDA Memorandum, July 26, 1994: Recommendation for Deferral of Donors for Malaria Risk.
7. FDA Memorandum, July 28, 1993: Deferral of Blood and Plasma Donors Based on Medications.
8. FDA Memorandum, October 7, 1988: Revised Guidelines for the Collection of Platelets, Pheresis.
9. Guidance for Industry: Revised Precautionary Measures to Reduce the Possible Risk of Creutzfeldt-Jakob Disease (CJD) and New Variant Creutzfeldt-Jakob Disease (nvCJD) by Blood and Blood Products. November 23, 1999.
10. FDA Memorandum, April 23, 1992: Revised Recommendations for Testing Whole Blood, Blood Components, Source Plasma and Source Leukocytes for Antibody to Hepatitis C Virus Encoded Antigen (Anti-HCV).

11. FDA Memorandum, December 12, 1991: Clarification of FDA Recommendations for Donor Deferral and Product Distribution Based on the Results of Syphilis Testing.
  12. FDA Memorandum, December 11, 1996: Interim Recommendations for Deferral of Donors at Increased Risk for HIV-1 Group O Infection.
  13. FDA Accepts AABB Changes to HIV-1 Group O Donor Questions. Association Bulletin #97-5, August 1, 1997.
  14. FDA Memorandum, June 8, 1995: Recommendations for the Deferral of Current and Recent Inmates of Correctional Institutions as Donors of Whole Blood, Blood Components, Source Leukocytes, and Source Plasma.
- 
-



## Abbreviated Donor History Questionnaire

|  | Yes                      | No                       |                                      |
|--|--------------------------|--------------------------|--------------------------------------|
| 1. Are you feeling healthy and well today?   | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 2. Have you read the educational materials and had your questions answered?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| <b>In the past 48 hours</b>  |                          |                          |                                      |
| 3. Have you taken aspirin or anything that has aspirin in it?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| <b>In the past 6 weeks</b>   |                          |                          |                                      |
| 4. Female donors: Have you been pregnant or are you pregnant now? (Males: check "I am male.")  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> I am male   |
| <b>In the past 8 weeks have you</b>  |                          |                          |                                      |
| 5. Donated blood, platelets or plasma?   | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 6. Had any vaccinations or other shots?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 7. Had close contact with the smallpox vaccination site of someone else?   | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| <b>In the past 16 weeks</b>  |                          |                          |                                      |
| 8. Have you donated a double unit of red cells using an apheresis machine?   | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| <b>Since your last donation have you</b>   |                          |                          |                                      |
| 9. Had any new medical problems or diagnoses?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 10. Had any new medical treatments?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 11. Taken any of the medications on the Medication Deferral List?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 12. Been outside the United States or Canada?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 13. Come into contact with someone else's blood?   | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 14. Had an accidental needle-stick?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 15. Had sexual contact with anyone who has HIV/AIDS or has had a positive test for the HIV/AIDS virus?                                     | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 16. Had sexual contact with a prostitute or anyone else who takes money or drugs or other payment for sex?                                 | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 17. Had sexual contact with anyone who has ever used needles to take drugs or steroids, or anything <u>not</u> prescribed by their doctor? | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 18. Had sexual contact with anyone who has hemophilia or has used clotting factor concentrates?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 19. Female donors: had sexual contact with a male who has ever had sexual contact with another male? (Males: check "I am male.")           | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> I am male   |
| 20. Had sexual contact with anyone who was born in or lived in Africa?   | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 21. Come into contact with blood from a person who has hepatitis?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 22. Had sexual contact with a person who has hepatitis?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 23. Lived with a person who has hepatitis?   | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 24. Received money, drugs, or other payment for sex?   | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 25. Male donors: had sexual contact with another male, even once? (Females: check "I am female.")  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> I am female |
| 26. Had a tattoo?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 27. Had ear or body piercing?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 28. Been in juvenile detention, lockup, jail, or prison for more than 72 hours?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 29. Used needles to take drugs, steroids, or anything <u>not</u> prescribed by your doctor?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |
| 30. Have any of your relatives had Creutzfeldt-Jakob disease?  | <input type="checkbox"/> | <input type="checkbox"/> |                                      |



American Red Cross

Together, we can save a life

HOME NEWS SERVICES PRESS ROOM FAQs JOBS PUBLICATIONS MUSEUM

What to Expect When Donating Blood

Apheresis

What Happens to Every Blood Donation  
Blood Services Regions

Eligibility Guidelines

FAQs

HELP NOW

Donate Now

Give Blood

Tissue Donation

Volunteer

Planned Giving

Donate Goods

Find Your Local Red Cross

Enter Zip Code Here:

FIND

Or Browse Through A List of Chapters!

WEBSITES

Search Our Site

GO

Receive monthly news, tips and



Home English > Services > Biomedical Services > Blood Services > Eligibility Guidelines

BLOOD DONATION ELIGIBILITY GUIDELINES

**Note to users:** This list is not complete. Medical professionals are available at each blood collection center and details of each donor's health and activities are discussed in a confidential setting prior to blood donation. The final determination of eligibility is made at that time. Some donor eligibility rules are specified by the Food and Drug Administration for every blood bank in the country. Other rules are determined by the particular blood bank and may differ between programs. Donor eligibility rules are intended to protect the health and safety of the donor as well as the patient who will receive the transfusion. The criteria listed below are provided as guidelines to assist you in determining whether you may be eligible to be a blood donor. The guidelines listed below were last revised on 6/27/02. There may have been some changes to these criteria since the last revision date. The most up to date eligibility information can be obtained by contacting the American Red Cross blood center nearest you.

GENERAL GUIDELINES

To give blood for transfusion to another person, you must be healthy, be at least 17 years old, weigh at least 110 pounds, and not have donated blood in the last 56 days. "Healthy" means that you feel well and can perform normal activities. If you have a chronic condition such as diabetes or high blood pressure, "healthy" also means that you are being treated and the condition is under control.

Other aspects of each potential donor's health history are discussed as part of the donation process before any blood is collected. Each donor receives a brief examination during which temperature, pulse, blood pressure and blood count (hemoglobin or hematocrit) are measured.

Making donations for your own use during surgery (autologous blood donation) is considered a medical procedure and the rules for eligibility are less strict than for regular volunteer donations.

Accupuncture  
Age

Hepatitis, Jaundice  
Hepatitis Exposure

updates from the Red Cross.

Email:

**SUBMIT**

#### OUR SUPPORTERS

► Major Gifts  
Provide Essential  
Support of Red  
Cross Disaster  
Relief Fund

► GE Foundation  
Donates \$1  
Million For Red  
Cross Disaster  
Relief

► Duracell and  
Safe Families

► Look Who's  
Helping

► Annual Disaster  
Giving Program

► Nextel Golf  
Tournament  
Benefits Red  
Cross

**CapitalOne**

Reach out  
to your community



[Click here for more info](#)

Allergy, Stuffy Nose, Itchy Eyes, Dry Cough  
Antibiotics  
Aspirin  
Asthma  
Autoimmune Diseases  
Birth Control  
Blood Pressure, High  
Blood Pressure, Low  
Blood Transfusion  
Cancer  
Cholesterol, high  
Chronic Illnesses  
Clotting Disorders  
Cocaine  
Cold, Flu, Sore Throat  
Creutzfeldt-Jakob Disease (CJD)  
Creutzfeldt-Jakob Disease, Variant (vCJD);  
"Mad Cow Disease"  
Dental Procedures  
Depression, Anxiety  
Diabetes mellitus  
Donation Intervals  
Epilepsy, Seizures  
Heart Disease  
Heart Murmur, Heart Valve Disorder  
Hemochromatosis  
Hemoglobin, Hematocrit, Blood Count

Herpes  
HIV, AIDS  
Hormone Replacement Therapy (HRP)  
Human Papilloma Virus (HPV)  
Hypertension, High Blood Pressure  
Immunization, Vaccination  
Infections  
Infectious Mononucleosis, "Mono"  
Intravenous Drug Use  
Lyme disease  
Malaria  
Marijuana, "Weed", "Pot"  
Medications  
Menstruation  
Organ/Tissue Transplants  
Piercing (ears, body), Electrolysis  
Pregnancy, Nursing  
Sexually Transmitted Disease  
Sickle Cell  
Skin Disease, Rash, Acne  
Surgery  
Syphilis/Gonorrhea  
Tattoo  
Tuberculosis  
Travel Outside of U.S., Immigration  
Vaccinations  
Venereal Diseases  
Weight

#### Accupuncture

Donors who have undergone acupuncture treatments are acceptable as long as the donor can confirm that the needles used in the treatment were sterile. Donors who cannot confirm that sterile needles were used in the acupuncture treatment are deferred from donating for 12 months.

[Back to Top](#)

#### Age

You must be at least 17 years old to donate to the general blood supply. Learn more about the reasons for a lower age limit. There is no upper age limit for blood donation as long as you are well with no restrictions or limitations to your activities.

[Back to Top](#)

**Allergy, Stuffy Nose, Itchy Eyes, Dry Cough**  
Acceptable as long as you feel well, have no fever, and have no problems breathing through your mouth.

[Back to Top](#)

**Antibiotics**  
Wait 2 days after finishing antibiotics for an infection (bacterial or viral). Acceptable if you are taking antibiotics to prevent an infection, for example, following dental procedures or for acne. Antibiotics for acne do not disqualify you from donating. If you have a temperature above 99.5 F, you may not donate until the fever is passed.

[Back to Top](#)

**Aspirin**  
See "Medications"

[Back to Top](#)

**Asthma**  
Acceptable as long as you are not having difficulty breathing at the time of donation and you otherwise feel well. Medications for asthma do not disqualify you from donating.

[Back to Top](#)

**Autoimmune Diseases**  
You are not eligible to donate if you have some types of generalized autoimmune disease including systemic lupus erythematosus and multiple sclerosis.

[Back to Top](#)

**Birth Control**  
Women taking birth control pills are acceptable.

[Back to Top](#)

**Blood Pressure, High**  
Acceptable as long as your blood pressure is below 180 systolic (first number) and below 100 diastolic (second number) at the time of donation. Medications for high blood pressure do not disqualify you from donating.

[Back to Top](#)

**Blood Pressure, Low**

Acceptable as long as you feel well when you come to donate. If your blood pressure normally runs low, it may be more difficult for your body to adjust to the volume loss following donation, especially if you are dehydrated. Drinking extra water before and after donation is important.

[Back to Top](#)

**Blood Transfusion**

Wait for 12 months after receiving a blood transfusion from another person in the United States. You may not donate if you received a transfusion since 1980 in the United Kingdom (England, Wales, Scotland, Northern Ireland, Channel Islands, Isle of Man), Gibraltar or Falkland Islands. This requirement is related to concerns about variant CJD, or 'mad cow' disease. Learn more about variant CJD and blood donation.

[Back to Top](#)

**Cancer**

Acceptable if the cancer was treated with only surgery or radiation, and it has been at least 5 years since treatment was completed with no cancer recurrence. If your cancer was treated with chemotherapy, hormonal therapy or immunotherapy, you are not eligible to donate. If you had leukemia or lymphoma, including Hodgkins Disease, you are not eligible to donate. Some low-risk cancers including squamous or basal cell cancers of the skin do not require a 5 year waiting period.

Precancerous conditions of the uterine cervix do not disqualify you from donation if the abnormality has been treated successfully.

You should discuss your particular situation with the health historian at the time of donation.

[Back to Top](#)

**Cholesterol, high**

Acceptable. Medications to lower the cholesterol level do not disqualify you from donating.

[Back to Top](#)

**Chronic Illnesses**

Most chronic illnesses are acceptable as long as you feel well, the condition is under good control, you have an adequate hemoglobin level and your temperature is normal when you come to donate. Chronic fatigue syndrome, fibromyalgia, rheumatoid arthritis, hypothyroidism, ulcerative colitis and Crohn's Disease do not automatically disqualify you from donating. You should discuss your condition with the health historian at the time of donation.

[Back to Top](#)

**Clotting Disorders**

If your blood does not clot normally, you should not donate since you may have excessive bleeding where the needle was placed. For the same reason, if you are taking any "blood thinner" (such as coumadin or heparin) you should not donate. If you are on aspirin, it is OK to donate blood. However, you must be off of aspirin for at least 36 hours in order donate platelets by apheresis.

[Back to Top](#)

**Cocaine**

Wait 12 months after using cocaine or other street drugs through your nose before attempting to donate blood. This requirement is related to concerns about hepatitis and HIV. Learn more about hepatitis and blood donation.

[Back to Top](#)

**Cold, Flu, Sore Throat**

Wait if you have a fever or a productive cough (bringing up phlegm)

Wait if you feel unwell on the day of donation.

Wait 2 days after you have completed antibiotic treatment for sinus, throat or lung infection.

[Back to Top](#)

**Creutzfeldt-Jakob Disease (CJD)**

If you ever received a corneal (eye) transplant, a dura mater (brain covering) transplant or human pituitary growth hormone, you are not eligible to donate. Those who have a close blood relative who had Creutzfeldt-Jacob disease or who is in a family that has been told they have a genetic risk for Creutzfeldt-Jacob disease are also not eligible to donate. Learn more about CJD.

[Back to Top](#)

**Creutzfeldt-Jakob Disease, Variant (vCJD); "Mad Cow Disease"**

See under Travel Outside of U.S. Learn more about vCJD and blood donation.

[Back to Top](#)

**Dental Procedures**

Acceptable after teeth cleaning, scaling, root canal, fillings and tooth extractions as long as there is no infection present.

Wait for 3 days after having other types of oral surgery, or after treatment for an abscess or infection in the mouth.

Wait 2 days after finishing antibiotics for a dental infection.

[Back to Top](#)**Depression, Anxiety**

Acceptable as long as you feel well and comfortable with the blood donation process. Medications for depression or anxiety do not disqualify you from donating.

[Back to Top](#)**Diabetes mellitus**

Acceptable two weeks after starting insulin. Medications to lower your glucose level do not disqualify you from donating. Those who since 1980, received an injection of bovine (beef) insulin made from cattle from the United Kingdom are not eligible to donate. This requirement is related to concerns about variant CJD, or 'mad cow' disease. Learn more about variant CJD and blood donation.

[Back to Top](#)**Donation Intervals**

Wait at least 8 weeks between whole blood (standard) donations.  
Wait at least 3 days between plateletpheresis donations.  
Wait at least 16 weeks between double red cell (automated) donations.

[Back to Top](#)**Epilepsy, Seizures**

Acceptable as long as you have been seizure-free for the last 3 months. Medications for seizures do not disqualify you from donating.

[Back to Top](#)**Heart Disease**

In general, acceptable as long as you have no restrictions on your physical activities, take no medications for heart disease other than aspirin, and have no current (within the last 6 months) heart-related symptoms such as chest pain.  
Wait at least 6 months following an episode of angina.  
Wait at least 6 months following a heart attack.  
Wait at least 6 months after bypass surgery or angioplasty.  
If you have a pacemaker, you may donate as long as your pulse is between 50 and 100 beats per minute with no more than a small number of irregular beats, and you meet the other heart disease criteria. You should discuss your particular situation with the health historian at the time of donation.

[Back to Top](#)

**Heart Murmur, Heart Valve Disorder**

Acceptable if you have a heart murmur as long as you have not had symptoms in the last 6 months, have no restrictions on your physical activity and are not taking any medications for heart disease other than prophylactic antibiotics (to prevent infections) or aspirin.

[Back to Top](#)

**Hemochromatosis**

American Red Cross does not accept individuals with hemochromatosis as blood donors for other persons at this time. Red Cross will continually re-evaluate this policy as more information accumulates.

[Back to Top](#)

**Hemoglobin, Hematocrit, Blood Count**

Acceptable if you have a hemoglobin at or above 12.5 g/dL.  
Acceptable if you have a hematocrit at or above 38%.

[Back to Top](#)

**Hepatitis, Jaundice**

If you had hepatitis (inflammation of the liver) caused by a virus, or unexplained jaundice (yellow discoloration of the skin), since age 11, you are not eligible to donate blood. This includes those who had hepatitis with infectious mononucleosis.

Acceptable if you had jaundice or hepatitis caused by something other than a viral infection, for example: medications, Gilbert's disease, bile duct obstruction, alcohol, gallstones or trauma to the liver.

If you ever tested positive for hepatitis B or hepatitis C, at any age, you are not eligible to donate, even if you were never sick or jaundiced from the infection.

[Back to Top](#)

**Hepatitis Exposure**

Wait 12 months after close contact with someone who is sick with viral hepatitis. Close contact is defined as sexual contact or sharing the same household, kitchen, dormitory, or toilet facilities.

Wait 12 months after detention in a correctional institution or residence in a long-term psychiatric institution.

Wait 12 months after receiving a blood transfusion (unless it was your own "autologous"



blood, blood injections, tattoo, non-sterile needle stick/body piercing or exposure to someone else's blood.

Wait 12 months following a human bite, if it broke the skin.

Wait 12 months after using cocaine or other street drugs through your nose.

Back to Top

### Herpes

Acceptable as long as you are feeling well.

Back to Top

### HIV, AIDS

Those who are at increased risk for becoming infected with HIV are not eligible to donate blood. According to the Food and Drug Administration, you are at increased risk if:

- you are a male who has had sex with another male since 1977, even once;
- you have ever used a needle, even once, to take drugs or steroids that were not prescribed by a physician;
- you have taken clotting factor concentrates for a bleeding disorder such as hemophilia;
- you were born in or lived in Cameroon, Central African Republic, Chad, Congo, Equatorial Guinea, Gabon, Niger, or Nigeria since 1977 (This requirement is related to concerns about HIV Group O. Learn more about HIV Group O.)
- you have taken drugs or money in exchange for sex since 1977;
- you have ever had a positive test for HIV virus;
- you have symptoms of HIV infection including unexplained weight loss, night sweats, blue or purple spots on or under the skin, long-lasting white spots or unusual sores in your mouth, lumps in your neck, armpits, or groin that last more than a month, fever higher than 99 degrees that lasts more than 10 days, diarrhea lasting over a month, or persistent cough and shortness of breath;

Wait for 12 months after close contact with someone who is at an increased risk for HIV infection. This occurs when paying to have sex, as a result of rape, or when having sex with an IV drug user.

Back to Top

### Hormone Replacement Therapy (HRP)

Women on hormone replacement therapy for menopausal symptoms and prevention of osteoporosis are eligible to donate.

Back to Top

**Human Papilloma Virus (HPV)**  
See "Venereal Diseases"

[Back to Top](#)

**Hypertension, High Blood Pressure**  
See "Blood Pressure, High

[Back to Top](#)

**Immunization, Vaccination**  
Acceptable if you were vaccinated for influenza, tetanus or meningitis, providing you are symptom-free and fever-free.

Wait 4 weeks after immunizations for German Measles (Rubella), MMR (Measles, Mumps and Rubella) and Chicken Pox.

Wait 2 weeks after immunizations for Red Measles (Rubeola), Mumps, Polio (by mouth), and Yellow Fever vaccine.

Wait 7 days after immunization for Hepatitis B as long as you are not given the immunization for exposure to hepatitis B.

- **Smallpox vaccination and did not develop complications**  
Wait 2 months (60 days) from the date of having a smallpox vaccination as long as you have had no complications. Complications may include skin reactions beyond the vaccination site or general illness related to the vaccination.
- **Smallpox vaccination and developed complications**  
Wait 14 days after all vaccine complications have resolved or 2 months (60 days) from the date of having had the smallpox vaccination whichever is the longer period of time. You should discuss your particular situation with the health historian at the time of donation. Complications may include skin reactions beyond the vaccination site or general illness related to the vaccination.
- **Smallpox vaccination – close contact with someone who has had the smallpox vaccine in the last eight weeks and you did not develop any skin lesions or other symptoms.**  
Eligible to donate.
- **Smallpox vaccination – close contact with someone who has had the vaccine in the last eight weeks and you have since a developed localized skin lesion only**  
Wait 2 months (60 days) from the date of the first skin lesion or sore. You should discuss your particular situation with the health historian at the time of donation. Complications may include skin reactions or general illness related to the exposure.
- **Smallpox vaccination – close contact with someone who has had the vaccine in the last eight weeks and you have since developed localized skin lesions and other complications**

Wait 14 days after all vaccine complications have resolved or 2 months (60 days) from the date of the first symptom whichever is the longer period of time. You should discuss your particular situation with the health historian at the time of donation. Complications may include skin reactions or general illness related to the exposure.

[Back to Top](#)

#### **Infections**

If you have a fever or an active infection, wait until the infection has passed before donating blood.

Wait 2 days after finishing antibiotics for an infection (bacterial or viral).

Infections with common herpes virus (cold sores or genital herpes) and Human Papilloma Virus (HPV) are acceptable as long as you feel well and do not have a fever.

Those who have had infections with Chagas Disease, babesiosis or leishmaniasis are not eligible to donate blood.

See also Antibiotics, Infectious Mononucleosis, Hepatitis, HIV, Syphilis/Gonorrhea, and Tuberculosis.

[Back to Top](#)

#### **Infectious Mononucleosis, "Mono"**

Acceptable if you had infectious mononucleosis ("mono") once the infection has passed, as long as you did not have hepatitis with the mononucleosis.

[Back to Top](#)

#### **Intravenous Drug Use**

Those who have ever used IV drugs that were not prescribed by a physician are not eligible to donate. This requirement is related to concerns about hepatitis and HIV. Learn more about hepatitis and blood donation.

[Back to Top](#)

#### **Lyme disease**

If this is a chronic condition you cannot donate. If you were treated with antibiotics and completely recovered, you can donate 12 months after the last dose of antibiotics was taken.

[Back to Top](#)

**Malaria**

Wait 3 years after completing treatment for malaria. Wait 12 months after returning from a trip to an area where malaria is found. Wait 3 years after moving to the United States after living in a country where malaria is found. Learn more about malaria and blood donation.

[Back to Top](#)

**Marijuana, "Weed", "Pot", "Ganga"**

Acceptable as long as you are not under the influence of marijuana at the time of donation.

[Back to Top](#)

**Medications** In almost all cases, medications will not disqualify you as a blood donor. Your eligibility will be based on the reason that the medication was prescribed. As long as the condition is under control and you are healthy, blood donation is usually permitted.

There are a handful of drugs that are of special significance in blood donation. Persons on these drugs have waiting periods following their last dose before they can donate blood:

- Accutane (isoretinoin), Proscar (finasteride), and Propecia (finasteride) - wait 4 weeks.
- Arava (leflunomide) - wait 3 months
- Avodart (dutasteride) - wait 6 months from the last dose
- Aspirin, no waiting period for donating blood. However you must wait 36 hours after taking aspirin or any medication containing aspirin before donating platelets by apheresis
- Chemotherapy-type drugs used for conditions other than cancer ( examples: bleomycin, interferon, methotrexate) - wait 4 weeks from last dose
- Coumadin, heparin or other prescription blood thinners- you should not donate since your blood will not clot normally. If your doctor discontinues your treatment with blood thinners, wait 5 days before returning to donate.
- human pituitary-derived growth hormone at any time - you are not eligible to donate blood
- Lupron used for condition other than cancer - wait 4 months from last dose
- Plavix - wait 36 hours after taking this medication before donating platelets by apheresis
- Soriatane (acitretin) - wait 3 years
- Tegison (etretinate) at any time - you are not eligible to donate blood
- Ticlid - wait 36 hours after taking this medication before donating platelets by apheresis

If you ever took Tegison (etretinate), you are not eligible to donate blood. If you ever took human pituitary-derived growth hormone, you are not eligible to donate blood. If you take aspirin, you can donate blood. However you must wait 36 hours after taking aspirin or any medication containing aspirin before donating platelets by apheresis. If you take Ticlid or Plavix, wait 36 hours after taking these medications before donating platelets by apheresis. If you are taking prescription blood thinners such as Coumadin or heparin, you should not

donate since your blood will not clot normally. If your doctor discontinues your treatment with blood thinners, wait 5 days before returning to donate.

[Back to Top](#)

#### **Menstruation**

Women may donate during their period if feeling well on the day of donation.

[Back to Top](#)

#### **Organ/Tissue Transplants**

Wait 12 months after receiving an organ or tissue transplant from another person. This includes bone and dental powder. If you are taking medications to prevent rejection of the organ or tissue you are not eligible to donate.

If you ever received a corneal (eye) transplant or a dura mater (brain covering) transplant, you are not eligible to donate. This requirement is related to concerns about the brain disease, Creutzfeld-Jacob Disease (CJD). Learn more about CJD and blood donation.

[Back to Top](#)

#### **Piercing (ears, body), Electrolysis**

Acceptable as long as the instruments used were sterile.

Wait 12 months if there is any question whether or not the instruments used were sterile and free of blood contamination. This requirement is related to concerns about hepatitis. Learn more about hepatitis and blood donation.

[Back to Top](#)

#### **Pregnancy, Nursing**

Persons who are pregnant are not eligible to donate. Wait 6 weeks after giving birth. Acceptable if you are nursing, or recently had an elective abortion.

[Back to Top](#)

#### **Sexually Transmitted Disease**

Wait 12 months after treatment for syphilis or gonorrhea.

[Back to Top](#)

#### **Sickle Cell**

Acceptable if you have sickle cell trait. Those with sickle cell disease are not eligible to donate.

[Back to Top](#)

#### **Skin Disease, Rash, Acne**

Acceptable as long as the skin over the vein to be used to collect blood is not affected. If the skin disease has become infected, wait until the infection has cleared before donating. Taking antibiotics to control acne does not disqualify you from donating.

[Back to Top](#)

#### **Surgery**

Acceptable once the wound is healed and stitches are dissolved or removed, as long as the underlying condition is also acceptable in a blood donor. Wait 2 days after having stitches or staples for lacerations. If a laceration has become infected, wait until the infection has cleared before donating. Wait 12 months if you had a blood transfusion from another person during surgery.

[Back to Top](#)

#### **Syphilis/Gonorrhea**

Wait 12 months after being treated for syphilis or gonorrhea.

[Back to Top](#)

#### **Tattoo**

Wait 12 months after a tattoo. This requirement is related to concerns about hepatitis. Learn more about hepatitis and blood donation.

[Back to Top](#)

#### **Tuberculosis**

Acceptable if you have a positive skin test for tuberculosis, or if you are receiving antibiotics for a positive TB skin test only. If you are being treated for a tuberculosis infection, wait until treatment is successfully completed before donating.

[Back to Top](#)

#### **Travel Outside of U.S., Immigration**

Wait 12 months after travel in an area where malaria is found. Wait 3 years after moving to the United States after living in a country where malaria is found. Persons who have spent long periods of time in countries where "mad cow disease" is found are not eligible to donate.

This requirement is related to concerns about variant Creutzfeldt Jacob Disease (vCJD). Learn more about vCJD and donation. Persons who were born in or who lived in certain countries in Western Africa, or who have had close contact with persons who were born in or who lived in certain West African countries are not eligible to donate. This requirement is related to concerns about HIV Group O. Learn more about HIV Group O, and the specific African countries where it is found.

[Back to Top](#)

#### Venereal Diseases

See also "Sexually Transmitted Disease"  
Wait 12 months after treatment for syphilis or gonorrhea.

[Back to Top](#)

#### Weight

You must weigh at least 110 Lbs to be eligible for blood donation for your own safety. Blood volume is in proportion to body weight. Donors who weigh less than 110Lbs may not tolerate the removal of the required volume of blood as well as those who weigh more than 110Lbs. There is no upper weight limit as long as your weight is not higher than the weight limit of the donor bed/lounge you are using. You can discuss any upper weight limitations of beds and lounges with your local health historian.

[Back to Top](#)

*Last updated: 6/27/02*

*By: R.A.R.,MD and M.A.P., RN,BSN*

***Note to users:** Eligibility guidelines may have changed since this information was last updated. For current information, please contact the American Red Cross blood region nearest you.*


© Copyright 2003 The American National Red Cross. All Rights Reserved.    [CONTACT US](#) | [SITE DIRECTORY](#) | [PRIVACY POLICY](#)

[home](#)
[about the tour](#)
[about blood](#)
[being a donor](#)
[recipient stories](#)
[sponsor a drive](#)
[young blood](#)
[games & more](#)

[DONATE BLOOD NOW](#)

[Schedule an appointment to give blood now!](#)
[Go!](#)

[SAVE A LIFE TOUR 2003](#)
[Am rican Red Cross](#)



## Apheresis

In an apheresis (ay-fur-ee-sis) donation, from the Greek "to take away," donors give only select blood components — platelets, plasma, red cells, infection-fighting white cells called granulocytes, or a combination of these, depending on the donors' blood type and the needs of the community. Apheresis is most commonly used to collect platelets and plasma.

*"By donating blood, I can serve my country in a very easy way."*

— Latia

» Register with Us

» Top 10

» How To Help

» Blood Donation Process

» Apheresis

» Tips for a Good Donation Experience

» Donor Eligibility

» Safety

» FAQ

» Thank You, Donors!

### Patient Benefits

A single apheresis donation of platelets can provide as many platelets as 5 whole blood donations. In addition, a platelet transfusion from a single donor greatly reduces the chances of an immune system reaction to the transfusion. Bone marrow transplant, cancer and leukemia patients whose immune systems are already compromised, benefit particularly from single donor platelet transfusions.

### Whole Blood vs. Apheresis

Apheresis donors' donations go through additional typing called Human Leukocyte Antigen (HLA) typing to ensure that the match between donor and recipient is as close as possible. Donors are then matched with specific patients in hospitals. Apheresis donors may receive emergency requests to donate for a patient to whom they are matched. Many apheresis donors find the knowledge that they are helping a specific individual in need particularly rewarding.




People who donate just platelets can donate every 3 days for a maximum of 24 times a year.

### Who Can Be an Apheresis Donor?


The same good health requirements that govern whole blood donors apply to apheresis donors. You must be at least 17 years old, weigh at least 110 pounds and be in good health.

### The Apheresis Donation Process: Safe and Easy

Similar to a whole blood donation, an apheresis donation consists of four steps: registration, health history and mini-physical, donation, and refreshments. From registration to refreshments, the process








lasts 1½ -2½ hours. During the actual donation, you will sit in a comfortable recliner, and a carefully monitored machine will draw blood from one arm through sterile tubing into a cell separator centrifuge. The blood stays inside the self-contained sterile tubing and never comes in contact with the machine. After the blood component(s) have been collected, the rest of the blood is returned to the donor through the same arm or the other arm. It's a safe process — the collection sets and needles are sterile, used once for each donor and then discarded. Donors usually relax, read, or enjoy a movie during the donation.

#### **Signing up**

Apheresis donations are by appointment only — call 1-800-GIVE LIFE to schedule an appointment.

© Copyright The American Red Cross. All Rights Reserved.

[RedCross.org](http://RedCross.org) | [Red Cross Store](#) | [Privacy Policy](#) | [Blood Information](#) | [Contact Us](#) | [Search](#) | [Links](#)



**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Application of:

GOODNOW

Serial No: 09/616,283

Filed: July 14, 2000

For: SYSTEM FOR DETECTING  
BACTERIA IN BLOOD, BLOOD  
PRODUCTS, AND FLUIDS OF  
TISSUES

Art Unit: 1645

Attorney Docket No. VRXB-P01-001

Examiner: J. Hines

Assistant Commissioner of Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

**Declaration Under 37 C.F.R. §1.132**

Sir:


I, Stephen J. Wagner, Ph.D., of Columbia, Maryland, hereby declare as follows:

1. I am the Director of Cell Therapy at the Holland Laboratory of the American Red Cross.

I have been conducting research in the field of transfusion medicine for over 14 years.

Accordingly, my curriculum vitae is attached as Appendix A.

2. I have read the above-identified application, the pending claims, and the Office Action mailed on September 29, 2003.
3. I understand that the Examiner has stated that the invention as described and claimed in the above-identified application is obvious in view of the teachings of Fisher et al. WO 98/57994, McLaughlin (U.S. Patent 4,683,196), Tadler et al. (*J. Clin. Lab. Anal.* 3: 21-25 (1989)), Erich et al. (*J. Immunol.* 143(12): 4053-4060, 1989), and Chang et al. (U.S. Patent 5,200,323).



For the reasons stated below, I respectfully disagree with the Examiner. I have been working in the field of transfusion medicine for over fourteen (14) years. During this time, the issue of bacterial contamination has been a source of primary concern for blood collectors and transfusion service scientists and clinicians. Accordingly, a number of the nation's leading blood safety experts have repeatedly called for initiation of a program to detect the presence of bacteria in blood and blood products.

5. In general, development of an effective test for detecting bacterial contamination is a formidable task and is unlike the development of effective tests for various viral agents for two reasons: (1) a bacterial test should detect the presence of the whole organism rather than the donor's immune response because donors may already have antibodies from previous exposures to certain bacteria; and (2) in many instances transfusion associated sepsis involves non-pathogenic environmental bacterial contaminants as opposed to pathogenic contaminants. Furthermore, an ideal test should be able to detect all or most of the diverse bacterial species implicated in transfusion-associated sepsis. *See, Stephen J. Wagner, (Zentralblatt für Bacteriologie 283(3): 253-257 (1996).*
6. In particular, developing immunological-based test is further complicated because it was thought that there were likely to be no exposed common antigens on the surface of all the diverse bacterial species implicated in transfusion associated bacterial sepsis.
7. In fact, at least one company, Binax used an immunological-based approach and failed. Binax in collaboration with the American Red Cross, developed a prototype assay using immunological methods to detect bacterial contamination of blood components. The test was developed to detect the presence of microbial antigens which were exposed after

physical or chemical treatments and subsequently visualized by immunochromatography.

However, Binax subsequently abandoned the test due to a limitation in its ability to detect all the bacteria implicated in platelet transfusions at levels below  $10^5$  CFU/mL.

8. As of 2001, no practical test was available to rapidly and accurately test for clinically relevant amounts of bacteria using antibodies. (Seaver et al. *Transfusion* 41: 1351-1355 (2001)) To the best of my knowledge no such reliable test is currently on the market.
9. To date, to the best of my knowledge, despite experimental measures to sterilize the collection of blood, transfusion-associated bacterial sepsis continues to be a concern in transfusion medicine. Further, despite many attempts, no detection technique has been implemented that meets all the requirements for a successful rapid test to be used just prior to transfusion.
10. Due to the failure of other immunoassays to detect bacteria in blood with adequate sensitivity, investigators in the field of blood banking have questioned whether an immunoassay, such as the one described and claimed by Verax would be successful.
11. I believe that the assay developed by Verax may provide the blood banking industry with a long-awaited method to rapidly and effectively test for clinically relevant amounts of bacterial contaminants in blood, blood products, and tissues collected at donation centers. Because they have been able to detect clinically relevant levels of diverse species of organisms using their two antibody system where others have failed, I believe that the Verax test provides a non-obvious solution to meet this need in the blood banking arena.

12. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title XVIII of the United States Code and that willful false statements may jeopardize the validity of this Application for Patent or any patent issuing thereon.

Dated: 12/11/03

Signature: Stephen J. Wagner

Stephen J. Wagner

Title: Director of Cell Therapy

Holland Laboratory of the American Red Cross

**STEPHEN J. WAGNER**

American Red Cross Blood Services  
Jerome H. Holland Laboratory for the Biomedical Sciences

**Education**

B.S. Chemistry, University of Maryland, College Park, MD, 1977  
M.S. Biophysics, The Pennsylvania State University, University Park, PA, 1979  
Ph.D. Biophysics, The Pennsylvania State University, University Park, PA, 1981

**Experience**

1972-1977 Technician, Chemistry Dept., Univ. MD, College Park, MD  
1977 Technician, Jet Propulsion Lab., Pasadena, CA  
1977-1981 Teaching Assistant, Dept. of Biophysics, The Penn. State Univ., Univ. Park, PA  
1977-1981 Research Assistant, Dept of Biophysics, The Penn. State Univ., Univ. Park, PA  
1981-1982 Postdoctoral Fellow, Laboratory of Genetics and Recombinant DNA NCI-Frederick Cancer Research Facility, Frederick, MD  
1982-1986 R & D Scientist, Zetachron Inc., State College, PA  
1986-1989 Postdoctoral Scholar, Dept. of Molecular and Cell Biology, The Penn. State Univ. Univ. Park, PA  
1989-1992 Scientist I, Product Development Department, American Red Cross, Holland Laboratory for the Biomedical Sciences, Rockville, MD  
1992- 1997 Scientist II, Product Development Department, American Red Cross, Holland Laboratory for the Biomedical Sciences, Rockville, MD  
1997-2002 Senior Scientist, Product Development Department, American Red Cross, Holland Laboratory for the Biomedical Sciences, Rockville, MD  
2002-present Director Cell Therapy Development, Blood & Cell Therapy Development American Red Cross, Holland Laboratory for the Biomedical Sciences, Rockville, MD

**Honors, Awards, Professional Recognition**

Cum Laude, University of Maryland, College Park, 1977  
Phi Kappa Phi  
Alpha Chi Sigma Award, 1977, Univ. of MD; awarded to an outstanding senior graduating in chemistry  
The Daymon Runyon-Walter Winchell Fellowship Grant, 1982, NCI FCRC, Frederick, MD  
National Tiffany Award, 1997, from the American Red Cross for outstanding technical achievement  
Appointed to the editorial board of **TRANSFUSION** (2001-2003)  
Member of AABB Transfusion-Transmitted Diseases Committee (2001-present)  
Scientific Advisor to FDA Bacterial Contamination Working Group (2001-present)  
Ad Hoc Reviewer for SPIR and STTR Grants, Transfusion Medicine (2001-present)  
Member of the American Red Cross National IRB (1999-present)  
Member of the American Red Cross Patent Disclosure Committee (2001-present)

### **Grant Awards**

1994-1997, NIH NHLBI, HL-94-05-B, "Novel Photoactive Virucides for Red Cells and Platelets, Total Direct Costs \$700,526

2001-2005, NIH NHLBI, HL66779, "Virus photoinactivation – Dimethylmethylene blue in RBC, Total Direct Costs \$1,250,000

### **Professional Society Memberships**

American Society for Microbiology

American Society for Photobiology

American Association of Blood Banks

International Society for Blood Transfusion

American Society for Hematology

### **Academic appointment**

Adjunct Associate Professor, Bowling Green State University, 1993, research advisor to Jean T. Scherrer: Master of Science thesis: Comparison of bacterial growth in platelet concentrates prepared by the platelet-rich plasma and the buffy coat methods.

### **Publications**

Watson, RT, Machado, G, Conway B, Wagner SJ, Davis DD: A kinetic study of the reaction of OH radicals with a series of halogenated alkanes, Part II, J Phys Chem 81:256-262, 1977.

Ravishankara S, Wagner SJ, Fisher SD, Smith G, Schniff B, Watson RT, Tesi G, and Davis DD,: A kinetics study of the reaction of OH with aromatics and olefins, Int J Chem Kinetics 10: 783-804, 1978.

Wagner SJ, Keith A, Strong K, and Snipes W: A spin-label assay for metal ion chelation and complex formation, Anal Biochem 99:175-182, 1979.

Watson RT, Ravishankara S, Machado G, Wagner SJ, and Davis DD: A kinetics study of the temperature dependence of hydroxyl radical ( $\text{OH}[\text{}^2\text{I}]$ ) with 1,1-dichloroethane and with 1,2-dichloro-2,2-difluoro ethane, Int J Chem Kinetics 11:187-197, 1979.

Wagner SJ, Taylor W, Keith A, and Snipes W: Effects of acridine plus near UV light on *E. coli* membranes and DNA in-vivo, Photochem Photobiol 32:771-779, 1980.

Lewis R, Sander S, Wagner S, and Watson RT: A temperature dependent study of  $\text{Cl}(\text{}^2\text{P}_{3/2})$  with a series of simple alkanes, J Phys Chem 84:2009-2015, 1980.

Wagner S, Keith A, and Snipes W: Interaction of divalent cations and proteins with phospholipid vesicles, Biochem. Biophys. Acta 600:367-375, 1980

Wagner SJ, and Snipes W: Effects of acridine plus near-UV light on the outer membrane of *Escherichia coli*, Photochem Photobiol 36:255-258, 1981.

Wagner SJ, Feldman A, and Snipes W: Recovery from damage induced by acridine plus near-ultraviolet light in *Escherichia coli*, Photochem. Photobiol. 35:73-81, 1982

Wagner SJ, Stevens SE Jr., Nixon BT, Lambert DH, Quivey RG Jr., and Tabita FR: Nucleotide and deduced amino acid sequence of the *Rhodobacter sphaeroides* gene encoding the form II ribulose 1,5-bis-phosphate carboxylase and comparison with other form I and II sequences, FEMS Micro Letts 55:217-222, 1988.

Wagner, SJ, Stevens SE Jr., and Nixon BT: Isolation of *gln A* from the cyanobacterium PR-6, in Light Energy Transduction in Photosynthesis: Higher plant and bacterial models, (Stevens SE Jr., Bryant DA, eds.), Amer Soc Plant Physiology, pp. 151-153, 1988.

Wagner SJ, Friedman LI, and Dodd RY: Approaches to the reduction of Viral Infectivity in Cellular Blood Components and Single Donor Plasma. Trans Med Rev 5:18-32, 1991.

Dodd RY, Moroff G, Wagner S, Dabay MH Dorfman E, George V, Ribeiro A, Shumaker J, and Benade LE: Inactivation of viruses in platelet suspensions which retain their in vitro characteristics: Comparison of psoralen-UVA and merocyanine 540-visible light methods, Transfusion 31: 483-490, 1991.

Wagner SJ, Wagner KF, and Friedman LI: Virucidal levels of ozone induce hemolysis and hemoglobin degradation. Transfusion 31: 748-751, 1991

Moroff G, Wagner S, Benade L, and Dodd R: Factors influencing virus inactivation and retention of platelet properties following treatment with aminomethyltrimethylpsoralen and ultraviolet A light. Blood Cells 18:43-56, 1992.

Wagner SJ, Storry JR, Mallory DA, Stromberg RR, Benade LE, and Friedman LI: Red cell membrane alterations associated with virucidal methylene blue phototreatment. Transfusion 33:30-36, 1992.

Wagner SJ, White R, Wolf L, Chapman J, Robinette D, Lawlor TE, and Dodd RY: Determination of residual AMT and mutagenicity testing following PUVA treatment of platelet suspensions. Photochem. Photobiol. 57: 819-824, 1992

Wagner SJ, Thomas SP, Kaufman RI, Nixon BT, and Stevens SE: The *glnA* gene of the cyanobacterium *Agmenellum quadruplicatum* PR-6 is nonessential for ammonium assimilation. J. Bacteriol. 175:604-612, 1992.



Wagner SJ, Robinette D, and Dodd R: Factors affecting *Yersinia enterocolitica* (serotype O:8) viability in deliberately inoculated blood. *Transfusion* 33:713-716, 1993.

Wagner SJ, Bardossy L, Moroff G, Dodd RY, Blajchman MA: Assessment of the hemostatic effectiveness of human platelets treated with aminomethyltrimethyl psoralen (AMT) and ultraviolet a light (UVA) using a rabbit ear bleeding time technique. *Blood* 82:3489-3492, 1993.

Lytle CD, Wagner SJ, Prodouz KN: Antiviral activity of gilyvocarcin V plus UVA radiation. *Photochem. Photobiol.* 818-821, 1993.

Wagner SJ: Use of methylene blue an aminomethyltrimethyl psoralen to inactivate viruses in cellular blood components: A review of current progress. *Proc. 5<sup>th</sup> Hokkaido Symp Transf Med, Sapporo*, 1993, pp 88-94.

Wagner SJ, Robinette D, Storry J, Cheng XY, Shumaker J, Benade L: Differential sensitivity of viruses in red cells to methylene blue photosensitization. *Transfusion* 34:521-526, 1994.

Wagner SJ, Friedman I, Dodd RY: Transfusion-associated bacterial sepsis. *Clinical Microbiology Reviews* 7:290-302, 1994.

Friedman LI, Stromberg RR, Wagner SJ: Reducing the infectivity of blood components - what we have learned, In: *Blood supply: risks, perceptions and prospects for the future*, Nance ST, editor, American Association of Blood Banks, Bethesda, MD, 1994.

Abe H and SJ Wagner: Analysis of viral DNA, protein, and envelope damage after methylene blue, phthalocyanine derivative or Merocyanine 540 photosensitization. *Photochem Photobiol* 61,402-409, 1995.

Wagner SJ, MA Cifone, H Murli, RY Dodd, B Myhr: Mammalian genotoxicity assessment of methylene blue in plasma: implications for virus inactivation. *Transfusion* 35:407-413, 1995.

Wagner SJ, G Moroff, AJ Katz, LI Friedman: Comparison of bacteria growth in single and pooled platelet concentrates after deliberate inoculation and storage. *Transfusion* 35:298-302, 1995.

Wagner SJ, D Robinette, M Nazario, G Moroff: Bacterial levels in components prepared from deliberately inoculated whole blood held for 8 or 24 hours at 20-24°C. *Transfusion* 35:911-918, 1995.

Wagner SJ: The potential for bacterial testing of blood products. *Zentralblatt für Bakteriologie* 283:253-257, 1996.

Wagner SJ, Robbinette D: Evaluation of swirling, pH, and glucose for the detection of bacterial contamination in platelet concentrates. *Transfusion* 36:989-993, 1996.

Skripchenko A, D Robinette, SJ Wagner: Comparison of methylene blue and methylene violet for photoinactivation of intracellular and extracellular virus in red cell suspensions. *Photochem Photobiol* 65:451-455, 1997.

Abe H, Wagner SJ, Kuwabara M, Kamo N, Ikebuchi K, Sekiguchi S: Factors affecting M13 bacteriophage inactivation by methylene blue photosensitization. *Photochem Photobiol* 65:873-876, 1997.

Wagner SJ. Transfusion-related bacterial sepsis. *Curr Opin Hematol* 4:464-469, 1997.

Abe H, Ikebuchi K, Wagner SJ, Kuwabara M, Kamo N, Sekiguchi S: Potential involvement of both type I and type II mechanisms in M13 virus inactivation by methylene blue photosensitization. *Photochem Photobiol* 66:204-208, 1997.

Wagner, SJ, Leiby D, Burch JW: Transfusion-associated bacterial sepsis: A concise review. *Immunohematology* 14:33-35, 1998.

Wagner SJ, Skripchenko A, Robinette D, Foley JW, Cincotta L: Factors affecting virus photoinactivation by a series of phenothiazine dyes. *Photochem Photobiol* 67:343-349, 1998.

Wagner SJ, Skripchenko A, Robinette D, Mallory DA, Cincotta L: Preservation of red cell properties after virucidal phototreatment with dimethylmethylene blue. *Transfusion* 38:729-737, 1998.

Wagner S, Robinette: Evaluation of an automated microbiological blood culture device for detection of bacteria in platelet components. *Transfusion* 38:674-679, 1998.

Hirayama J, Wagner SJ, Gomez C, MacDonald V, Abe H, Ikeda H, Ikebuchi K, Sekiguchi S: Virus photoinactivation in stroma free hemoglobin (SFH) with methylene blue (MB) or 1,9-dimethylmethylene blue (DMMB). *Photochem Photobiol* 71:90-93, 2000 .

Haughtaling MA, Perera R, Owen KE, Wagner S, Kuhn RJ, Morrison H: Photobiological properties of positively charged methylene violet analogs. *Photochem Photobiol* 71:20-8, 2000.

Wagner SJ, Robinette D, Friedman LI, Miripol J: Diversion of initial blood flow to prevent whole blood contamination by "skin" surface bacteria: an in-vitro model. *Transfusion*, 40:335-338, 2000.

Wagner SJ, Skripchenko A, Robinette D, Mallory DA, Hirayama J, Cincotta L, Foley J: The use of dimethylmethylene blue for virus photoinactivation in red cell suspensions, in *Advances in Transfusion Safety*, Brown F, Vyas V, eds, Karger Press, Basel, 2000, pgs 125-130.

Skripchenko A, Wagner SJ: Photoinactivation of leukocytes in erythrocyte suspensions: Comparison of methylene blue and dimethylmethylene blue. *Transfusion*. 2000 Aug;40(8):968-75.

Hirayama J, Wagner SJ, Abe H, Ikebuchi K, Ikeda H. Involvement of reactive oxygen species in hemoglobin oxidation and virus inactivation by 1,9-dimethylmethylen blue phototreatment. *Biol Pharm Bull* 2001;24:418-421.

Wagner SJ, Skripchenko A, Pugh JC, Suchmann DB, Ijaz MK. Duck hepatitis B photoinactivation by dimethylmethylen blue in RBC suspensions. *Transfusion* 2001;41:1154-8.

Seaver, M, Crookston JC, Roselle, DC, Wagner, SJ. First results using automated epifluorescence microscopy to detect *Escheichia coli* and *Staphylococcus epidermidis* in leukoreduced platelet concentrates. *Transfusion* 2001 41: 1351-1355.

Wagner, SJ. Virus inactivation in blood components by photoactive phenothiazine dyes. *Tranfusion Medicine Reviews* 2002;16:61-66.

Wagner SJ. Pathogen Inactivation in blood components. In Rossi's principles in Transfusion Medicine, 3<sup>rd</sup> edition, Simon T, Senior Editor, Lippincott Williams & Wilkins, 2002, Philadelphia, PA, pp 806-814.

Wagner S, Skripchenko A, Thompson-Montgomery D. Use of a flow-cell system to investigate virucidal dimethylmethylen blue phototreatment in two RBC additive solutions. *Transfusion* 2002;42:1200-1205.

Wagner SJ, Skripchenko A, Thompson-Montgomery D. Quinacrine enhances vesicular stomatitis virus inactivation and diminishes hemolysis of dimethylmethylen blue phototreated cells. *Photochem Photobiol* 2002;76:514-517.

Wagner SJ, Myrup A, Xu Y-W, Katz AJ. Sterile isolation and cryopreservation of adherent monocytes/macrophages from mononuclear cell apheresis preparations. *Wound Rep Reg* 2003;11:145-49.

Wagner SJ. Pathogen reduction of bacteria and parasites, In: Bacterial and parasitic contamination of blood products, Brecher M, ed., AABB Press, Bethesda, MD, 2003.

## **Patents**

Snipes W, and Wagner SJ: Erodible matrix for sustained release bioactive composition; polyoxyethylene glycol, stearic acid or stearyl alcohol, Patent no. 4744976, 1988.

Snipes W, and Wagner SJ: Erodible matrix for sustained release bioactive composition: Solid water-dispersable polyether and erosion rate modifier, Patent no. 4806337, 1989.

Snipes W, and Wagner SJ: Controlled release potassium chloride composition: Capsules with fatty acids, Patent no. 4832955, 1989.

Wagner SJ: Inactivation of extracellular enveloped viruses in blood and blood components by

phenothiazin-5-ium dyes plus light, Patent no. 5545516, 1996.


Wagner SJ, Cincotta L: Intracellular and extracellular decontamination of whole blood and blood components by amphiphilic phenothiazine-5-ium dyes plus light. Patent no. 6030767, 2000.

Seaver M, Crookston JC, Roselle DC, Wagner SJ: Automated fluorescent microscopy for bacterial detection, in prosecution.

Wagner SJ, Myrup A, Celluzzi C: Method for enriching adherent monocyte populations, in prosecution.

Wagner SJ, Skripchenko A: Decontamination of biological fluids using diphenylpyrilium compounds, in prosecution.

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**



Re Application of:

GOODNOW

Serial No: 09/616,283

Filed: July 14, 2000

For: SYSTEM FOR DETECTING  
BACTERIA IN BLOOD, BLOOD  
PRODUCTS, AND FLUIDS OF  
TISSUES

Art Unit: 1645

Attorney Docket No. VRXB-P01-001

Examiner: J. Hines

Assistant Commissioner of Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

**Declaration Under 37 C.F.R. §1.132**

Sir:

I, Jeffrey A. Hall, Ph.D., of Franklin, MA, hereby declare as follows:

1. I am the Director of Assay Development at Verax Biomedical, Inc., the assignee of the present application. I have been conducting research in immunoassay development for 13 years. Accordingly, my curriculum vitae is attached as Appendix A.
2. I have read the above-identified application, the pending claims, the Office Action mailed on February 11, 2003 and the Office action mailed on September 29, 2003.
3. I understand that the Examiner has stated that the invention as described and claimed in the above-identified application is obvious in view of the teachings of Fisher et al. WO 98/57994, McLaughlin (U.S. Patent 4,683,196), Tadler et al. (*J. Clin. Lab. Anal.* 3: 21-25 (1989)), Erich et al. (*J. Immunol.* 143(12): 4053-4060, 1989), and Chang et al. (U.S. Patent 5,200,323).

4. The Examiner states that McLaughlin teaches antibodies which specifically bind to gram negative bacteria in order to determine their presence and/or absence while Tadler et al., teach well known binding agents that bind lipotechoic acid of gram-positive bacteria in assays. See Office Action dated February 11, 2003. In the current Office Action the Examiner admits that the McLaughlin antibodies are not pan-generic but states that it would have been to obvious modify the assay to include the antibodies taught by Erich et al. instead of the McLaughlin antibodies. Similarly, Examiner admits that the Tadler et al. antibodies are not pan-generic but states that it would have been obvious to modify the assay to include the antibodies taught by Fischer et al. instead of the Tadler et al. antibodies.
5. I have reviewed the disclosures of all the cited references: McLaughlin, Erich et al., Tadler et al., and Fischer et al. For the reasons set forth below and the accompanying experimental data, I believe that the prior art antibodies fail to demonstrate broad pan-generic cross-reactivity and detection at a level of sensitivity to be effective in detecting clinically relevant amounts of bacteria in a blood or blood products as required by the claims.
6. Verax Biomedical, Inc., has developed pan-generic antibodies immunoreactive with the Gram-negative antigen lipopolysaccharide (LPS) and pan-generic antibodies immunoreactive with the Gram-positive antigen lipoteichoic acid (LTA). The pan-generic activity and sensitivity of these antibodies has been compared with the closest commercially available antibodies that are being marketed for pan-generic reactivity. We

set forth below the comparative results obtained for the Verax gram positive as well as the gram negative antibodies.

### **Gram-Positive Antibodies**

7. The Examiner states that Tadler et al disclose well known binding agents that bind the lipoteichoic acid (LTA) of the Gram-positive bacteria. We purchased commercially available antibodies that bind the LTA of the Gram-positive bacteria [Appendix B ] and performed a side-by-side comparison with the Verax antibodies (VERAX PGD BA-3).<sup>1</sup>
8. The specification as filed discloses how to make and use pan-generic gram positive antibodies. Example 9 of the application demonstrates to one of ordinary skill in the art that the Verax monoclonal antibody clone 96-110 (described in WO 98/57994 by Fisher et al.; now designated as Verax PGD BA-3) shows pan-generic reactivity with seven Gram-positive bacteria as depicted in Figure 5 of the application. [Appendix C].
9. In addition, to the examples described in Example 9, we have generated and screened for additional antibodies that are pan-generic in nature and are capable of detecting clinically relevant amounts of bacteria using the methods taught in the specification at pages 24-26. VERAX PGD BA-4 is an example of such an antibody.
10. We conducted further assays comparing the Verax antibodies, such as VERAX PGD BA-3 and VERAX PGD BA-4 to commercially available antibodies from HyCult Biotech and Biogenesis, Inc. We evaluated the ability of these antibodies to detect clinically

---

<sup>1</sup> The VERAX PGD BA-3 antibody is the same as the antibody described in Example 9, i.e., the Fischer antibody of clone 96-110.

relevant amounts of bacteria and to be useful in constructing meaningful screening assays. These results are presented below:

## PAN-GENERA REACTIVITIES (S:N RATIO) OF VARIOUS BINDING AGENTS TOWARDS GRAM POSITIVE BACTERIA

|                 |         | TEST BACTERIA            |                              |                          |                          |                              |                              |                              |                            |                                |
|-----------------|---------|--------------------------|------------------------------|--------------------------|--------------------------|------------------------------|------------------------------|------------------------------|----------------------------|--------------------------------|
|                 |         | <i>Staph epidermidis</i> | <i>Staphylococcus aureus</i> | <i>Staph lugdenensis</i> | <i>Bacillus subtilis</i> | <i>Group B Streptococcus</i> | <i>Group G Streptococcus</i> | <i>Enterococcus faecalis</i> | <i>Corynebacterium sps</i> | <i>Clostridium perfringens</i> |
| Vendor          | Antigen |                          |                              |                          |                          |                              |                              |                              |                            |                                |
| HyCult Biotech  | G+ LTA  | 28.1                     | 1.2                          | 1.4                      | 1.3                      | 1.5                          | 3.8                          | 1.9                          | 3.8                        | 1.7                            |
| Biogenesis Inc. | G+ LTA  | 9.2                      | 16.3                         | 1.1                      | 2.9                      | 1.5                          | 6.7                          | 1.3                          | 6.7                        | 2.2                            |
| VERAX PGD BA-3  | G+ LTA  | 62.6                     | 7.6                          | 12.5                     | 20.3                     | 5.6                          | 14.8                         | 5.8                          | 14.8                       | 4.7                            |
| VERAX PGD BA-4  | G+ LTA  | 77.9                     | 30.1                         | 71.3                     | 10.2                     | 6.6                          | 10.8                         | 21.4                         | 10.7                       | 2.8                            |

\* "SAMPLE-TO-NOISE" RATIO = ANTIGEN-SPECIFIC SIGNAL/BACK-GROUND SIGNAL

\*\* S:N RATIO IS A COMMON EIA DATA NORMALIZATION TECHNIQUE TO SIMULTANEOUSLY COMPARE REACTIVITIES OF MULTIPLE BINDING AGENTS. A S:N RATIO >2 IS REQUIRED TO CONSTRUCT A MEANINGFUL ASSAY.

Our results show that the commercially available antibodies were not truly pan-generic with respect to the detection of the LTA on Gram-positive bacterium and would not be useful in constructing a meaningful blood screening assay. In contrast, the Verax antibodies detected seven genera of Gram-positive bacterium routinely found in contaminated blood. See results above and Figure 5.

11. To be effective, a signal: noise ratio greater than two (2) is required for constructing a meaningful assay. The commercially available antibodies failed to demonstrate a ratio greater than two. As can be seen from the Table above, the Hycult Biotech antibody



failed to detect two species of *Staphylococcus* bacterium, *Bacillus subtilis*, Group B *Streptococcus*, *Enterococcus*, *Corynebacterium*, and *Clostridium*. Similarly, the Biogenesis Inc. antibody failed to detect one species of *Staphylococcus* bacterium, Group B *Streptococcus* and *Enterococcus*. In contrast, the Verax antibodies were effective in detecting seven different genera of bacteria and the signal was considerably stronger than that observed for the commercially available antibodies.

12. We now provide a side-by-side comparison of the pan-generic reactivity of the Verax antibodies to the Tadler et al. antibodies. Tadler et al. discloses an immunoassay for the detection of the LTA on gram positive bacteria. A close review of their experimental data shows that their antibodies demonstrate limited binding and detection of four bacterial genera: *Streptococcus spp.*, *Staphylococcus spp.*, *Enterococcal spp.*, and *Clostridium*. In contrast, the Verax antibodies are capable of pan-generic binding and detection of at least seven Gram-positive bacterial genera.
13. We further provide a comparison of the sensitivity of the Verax antibodies to the Tadler et al. antibodies. Figure 2 of Tadler et al. shows that only 2 bacterial genera, i.e., *Streptococcus mutans* and *Staphylococcus epidermidis*, are detected at clinically relevant amounts,  $5 \times 10^5$  CFU/50 $\mu$ l (i.e.,  $1 \times 10^7$  CFU/ml). Thus, at this level of sensitivity the Tadler antibodies are not truly pan-generic. The bacteria *Staphylococcus aureus* is detected at  $5 \times 10^6$  CFU/50 $\mu$ l (i.e.,  $1 \times 10^8$  CFU/ml), a level that is not clinically relevant. Additionally, the Tadler et al. immunoassay was unable to detect *Staphylococcus faecium* at  $5 \times 10^6$  CFU/50 $\mu$ l ( $1 \times 10^8$  CFU/ml) suggesting that the antibodies are only able to cross-react with LTA on certain *Staphylococcus spp.* and *Streptococcus spp.* Therefore, in

effect, Tadler et al. show detection of only two genera of gram-positive bacteria at clinically relevant amounts as is required by the claims. See Figure 2 of Tadler et al.

14. In contrast, the following Table set forth both the pan-generic cross-reactivity of the Verax antibodies, as well as the sensitivity of these antibodies, demonstrating a greater degree of sensitivity in detecting clinically relevant amounts of bacteria in contaminated blood or blood products ( $1 \times 10^2$  CFU/ml –  $1 \times 10^6$  CFU/ml).

| GRAM POSITIVE RAPID TEST SIGNAL (G/DENS) |                       |                  |                       |                  |             |             |                    |                    |                        |                       |        |
|--|-----------------------|------------------|-----------------------|------------------|-------------|-------------|--------------------|--------------------|------------------------|-----------------------|--------|
| CFU/ml                                   | <i>S. epidermidis</i> | <i>S. aureus</i> | <i>S. lugdunensis</i> | <i>B. cereus</i> | GRP B Strep | GRP G Strep | <i>S. pyogenes</i> | <i>E. faecalis</i> | <i>C. minutissimum</i> | <i>C. perfringens</i> | CFU/ml |
| 1.0 E5                                   | 17.26                 | 1.04             | 13.21                 | 12.24            | 0.26        | 16.07       | nt                 | 13.95              | 6.82                   | 11.20                 | 1.0 E5 |
| 5.0 E4                                   | nt                    | nt               | 3.20                  | 9.91             | nt          | nt          | 21.77              | nt                 | nt                     | nt                    | 5.0 E4 |
| 1.0 E4                                   | 8.81                  | 0.41             | 0.36                  | 2.45             | 0.04        | 1.43        | 10.32              | 3.19               | 2.03                   | 8.67                  | 1.0 E4 |
| 5.0 E3                                   | 3.98                  | nt               | nt                    | nt               | 0.04        | nt          | nt                 | 0.44               | nt                     | nt                    | 5.0 E3 |
| 1.0 E3                                   | 0.83                  | 0.24             | 0.17                  | 0.29             | nt          | 0.46        | 0.51               | 0.27               | 1.26                   | 2.91                  | 1.0 E3 |

\*BOXED CELL = MINIMALL DETECTABLE CONCENTRATION

\*\*G/DENS = REFLECTANCE SIGNAL, ANY G/DENS > 0.25 IS VISIBLE

\*\*\*nt = NOT TESTED

15. We now examine the pan-generic cross-reactivity of the Fischer et al antibody as taught by the Fischer PCT publication . The Fischer PCT publication referenced by the Examiner fails to disclose or suggest pan-generic activity. The Table below provides a comparison of the properties shown in the Fischer references in comparison to the properties of the Fischer antibody now appreciated and recognized by Verax.

|  |  |
|--|--|
| Gram positive bacterial species recognized by the Fisher antibody as taught by the Fischer reference | Gram positive bacterial species recognized by the Fisher antibody as taught by the instant application |
|--|--|

|                            |  |
|----------------------------|--|
| <i>Staphylococcus spp.</i> | <i>Staphylococcus spp.</i><br><i>Streptococcus pyogenes</i><br><i>Group B Streptococcus</i><br><i>Group G Streptococcus</i><br><i>Enterococcus faecalis</i><br><i>Corynebacterium minutissimum</i><br><i>Clostridium perfringens</i><br><i>Bacillus spp.</i> |
|----------------------------|--|

Thus, the Fisher et al. reference shows binding to *only* one bacterial genus: the *Staphylococcus* genus: there was no recognition of its pan-generic ability in the Fischer reference.

#### **Gram-Negative Antibodies**

16. The Examiner states that McLaughlin and/or Erich et al. disclose antibodies which specifically bind to gram-negative bacteria. McLaughlin discloses mouse and rabbit antibodies that bind to the Lipid A core of the LPS on Gram-negative bacteria. Erich et al disclose three murine monoclonal antibodies and state that one of the three antibodies showed cross-reactivity with the heterologous LPS and Gram bacterial strains but cross-reactivity was either moderate or virtually absent for the remaining two. See Page 4058. We conducted a side-by-side comparison of the Verax antibodies to the closest commercially available mouse antibodies that are marketed as being anti-LPS core or anti-endotoxin [Appendix D] from HyCult Biotech, Virostat, and QED.
17. The specification as filed discloses how to make and use pan-generic gram negative antibodies. Example 9 of the application demonstrates to one of ordinary skill in the art

that the Verax monoclonal antibody clone 26-5 (commercially available from Biodesign International) shows pan-generic reactivity with the LPS of seven Gram-negative bacteria as depicted in Figure 6 of the application. [Appendix E].

18. In addition to the examples described in Example 9, we have generated and screened for additional antibodies that are pan-generic in nature and are capable of detecting clinically relevant amounts of bacteria using the methods taught in the specification at pages 24-26. VERAX PGD BA-1 and VERAX PGD BA-1 are examples of such antibodies.
19. McLaughlin antibodies are reactive with gram-negative bacteria such as *Neisseria*, *Chlamydia*, and *Salmonella*. The Erich et al antibodies are cross-reactive with only two (live) bacterial genera: *Escherichia* and *Salmonella*. In contrast, the Verax antibodies show pan-generic activity against a range of bacteria that have been identified as contaminants in blood and blood products in three major national transfusion reaction studies including the BaCon Study in the United States, the Hemovigilance study in France, and the SHOT study in the United Kingdom. These contaminants include both pathogenic as well as non-pathogenic bacterial species such as *Yersinia enterocolitica* and *Proteous miribilis*, along with other common soil-borne bugs. The McLaughlin and Erich et al antibodies appear to be effective mainly against the pathogenic gram negative bacterial species (e.g., *Neisseria*, *Escherichia*, and *Chlamydia*).
20. In addition, we conducted further assays using Verax antibodies, in evaluating the ability of these antibodies to detect clinically relevant amounts of bacteria and be useful in constructing meaningful screening assays. These results are presented below:

## PAN-GENERA REACTIVITIES (S:N RATIO) OF VARIOUS BINDING AGENTS TOWARDS GRAM NEGATIVE BACTERIA

|                |         | TEST BACTERIA               |                               |                                |                           |                              |                         |                               |                               |                                |                          |                            |
|----------------|---------|-----------------------------|-------------------------------|--------------------------------|---------------------------|------------------------------|-------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------|----------------------------|
|                |         | <i>Enterobacter cloacae</i> | <i>Enterobacter aerogenes</i> | <i>Acinetobacter baumannii</i> | <i>Klebsiella oxytoca</i> | <i>Klebsiella pneumoniae</i> | <i>Escherichia coli</i> | <i>Pseudomonas aeruginosa</i> | <i>Salmonella enteritidis</i> | <i>Yersinia enterocolitica</i> | <i>Proteus mirabilis</i> | <i>Serratia marcescens</i> |
| Vendor         | Antigen |                             |                               |                                |                           |                              |                         |                               |                               |                                |                          |                            |
| HyCult Biotech | LPS     | 1                           | 1                             | 1.1                            | 1.2                       | 1.1                          | 1.2                     | 1                             | 1                             | 1                              | 1.3                      | 1.3                        |
| Virostat       | LPS     | 1                           | 1                             | 1                              | 1                         | 1                            | 1                       | 1                             | 1                             | 1                              | 1                        | 1.1                        |
| QED            | LPS     | 1                           | 1                             | 1                              | 1                         | 1                            | 1                       | NT                            | NT                            | NT                             | NT                       | NT                         |
| VERAX PGD BA-1 | LPS     | 12.1                        | 11.1                          | 12.5                           | 12.8                      | 12.3                         | 9.6                     | 9.7                           | 11.2                          | 11.4                           | 10.6                     | 12.3                       |
| VERAX PGD BA-2 | LPS     | 11.2                        | 16.5                          | 14.2                           | 9.5                       | 7.8                          | 13.5                    | 9.2                           | 8.5                           | 12.1                           | 19.5                     | 22.7                       |

\* "SAMPLE-TO-NOISE" RATIO = ANTIGEN-SPECIFIC SIGNAL/BACK-GROUND SIGNAL

\*\* S:N RATIO IS A COMMON EIA DATA NORMALIZATION TECHNIQUE TO SIMULTANEOUSLY COMPARE REACTIVITIES OF MULTIPLE BINDING AGENTS  
A S:N RATIO >2 IS REQUIRED TO CONSTRUCT A MEANINGFUL ASSAY.


21. To be effective, a signal: noise ratio greater than two (2) is required for constructing a meaningful assay. As can be seen from the Table above, the commercially available antibodies failed to detect all nine bacterial genera. In contrast, the Verax antibodies were effective in detecting nine different genera of bacteria and showed strong signals for each of these bacteria. Thus, the Verax antibodies showed greater effectiveness than the commercially available antibodies.
22. In addition, the following Table sets forth the sensitivity of the Verax gram negative antibodies, demonstrating a greater degree of sensitivity in detecting clinically relevant amounts of bacteria in contaminated blood or blood products ( $1 \times 10^2$  CFU/ml –  $1 \times 10^6$  CFU/ml).

VERAX PLATELET PGD ASSAY: ANALYTICAL SENSITIVITY

| GRAM NEGATIVE RAPID TEST SIGNAL (G/DENS) |                |                      |                     |                   |                   |                     |                      |                       |                     |                      |        |
|--|----------------|----------------------|---------------------|-------------------|-------------------|---------------------|----------------------|-----------------------|---------------------|----------------------|--------|
| CFU/ml                                   | <i>E. coli</i> | <i>P. aeruginosa</i> | <i>E. aerogenes</i> | <i>K. oxytoca</i> | <i>E. cloacae</i> | <i>A. baumannii</i> | <i>K. pneumoniae</i> | <i>S. enteritidis</i> | <i>P. mirabilis</i> | <i>S. marcescens</i> | CFU/ml |
| 1.0 E5                                   | 4.51           | 2.16                 | 9.13                | 8.66              | 1.01              | 7.71                | 5.30                 | 1.14                  | 1.38                | 0.83                 | 1.0 E5 |
| 5.0 E4                                   | 4.10           | 1.45                 | 7.70                | 7.08              | 0.77              | 7.51                | 4.40                 | 0.70                  | 0.98                | 0.71                 | 5.0 E4 |
| 1.0 E4                                   | 2.75           | 0.79                 | 6.88                | 3.31              | 0.82              | 4.79                | 2.34                 | 0.50                  | 0.56                | 0.69                 | 1.0 E4 |
| 5.0 E3                                   | 1.34           | 0.67                 | 4.81                | 1.46              | 0.73              | 2.98                | 1.21                 | 0.48                  | 0.24                | 0.16                 | 5.0 E3 |
| 1.0 E3                                   | 0.99           | 0.32                 | 4.10                | 0.25              | 0.75              | 1.46                | 0.57                 | 0.04                  | 0.09                | 0.01                 | 1.0 E3 |

23. For these reasons, I believe that the closest commercially available prior art antibodies do not show the pan-generic activity or sensitivity required to be effective in detecting clinically relevant amounts of bacterial contaminants in blood and blood products.
24. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title XVIII of the United States Code and that willful false statements may jeopardize the validity of this Application for Patent or any patent issuing thereon.

Dated: 12/2/03

Signature:   
Dr. Jeffrey A. Hall, Ph.D  
Director, Assay Development  
Verax Biomedical Incorporated

## **CURRICULUM VITAE**

**Jeffrey A. Hall, Ph.D**

87 Skyline Dr.  
Franklin, MA. 02038  
(508)-541-8707 (home)  
(508)-755-7029 ex. 102 (work)  
email: jhall@veraxbiomedical.com  
email: jlehall@aol.com

### **DEGREES:**

- 1986-1988      Doctor of Philosophy (Reproductive Physiology), West Virginia University, Morgantown, West Virginia.
- 1982-1986      Master of Science (Reproductive Physiology), West Virginia University, Morgantown, West Virginia.
- 1978-1982      Bachelor of Science (Biological and Agricultural Sciences), Arizona State University, Tempe, Arizona.

### **WORK EXPERIENCE:**

- 2001-              **Director, Assay Development, Verax Biomedical Inc., Worcester, MA.**
- 1999 - 2000      **Manager, Industrialization Group - VIDAS Probe, bioMerieux, Inc., Rockland, MA.**
- 1997 - 1999      **Senior Staff Scientist, Infectious Disease Assay Development, Bayer Diagnostics (formerly Chiron Diagnostics Corp.), E. Walpole, MA.**
- 1995 -1997      **Staff Scientist, Infectious Disease Assay Development, Chiron Diagnostics Corp. (formerly Ciba-Corning Diagnostics), E. Walpole, MA.**
- 1994-1995      **Senior Scientist, Assay Development R&D, Dade International (formerly Baxter Diagnostics Inc.)**
- 1991-1994      **Scientist, Assay Development R&D, Baxter Diagnostics Inc. (Miami, FL)**
- 1988-1990      **Post-Doctoral Research Associate, Reproductive Physiology Unit, University of Missouri-Columbia (Columbia, MO.)**

### **HONOR SOCIETIES:**

Beta Beta Beta  
Gamma Sigma Delta  
Phi Kappa Phi  
Sigma Xi

## **AWARDS:**

- 1987 First Place Award, Sigma Xi Graduate Research Colloquium, West Virginia University  
1986-88 Swigert Award for Outstanding Doctoral Research Assistant, West Virginia University  
1985 Third Place Award, Northeast Section Am. Soc. Animal Science, Graduate Research Competition, West Virginia University

## **MEMBERSHIPS:**

American Association of Blood Bankers  
American Association of Clinical Chemistry  
American Society of Microbiology  
American Association for the Advancement of Science

## **REFEREED PUBLICATIONS:**

- Hall, J., W. Jilg, B. Hottentraeger, P. Bonnar, C. Fang, L. Baker. 1998. Performance of a chemiluminescent immunoassay for HBsAg on a new high-throughput and fully automated ACS:Centaur system. Clin. Lab. 44:349-354.
- Hall, J.A., R.A. Dailey, P.E. Lewis. 1993. Effect of the corpus luteum of pregnancy on ovulation and luteal function in early post-partum ewes. J. Anim. Sci. 71:3067-3072.
- Hall, J.A., B.N. Day, R.V. Anthony. 1993. Influence of steroids on relaxin-induced changes in distensibility and composition of the cervix in gilts. Biol. Reprod. 48:1348-1353.
- Hall, J.A., E.M. Meisterling, A.M. Benoit, D.A. Cooper, D.A. Coleman, S.P. Lerner, P.E. Lewis, R.A. Dailey. 1993. Factors contributing to the formation of experimentally-induced ovarian cysts in prepubertal gilts. Dom. Anim. Endocrinol. 102:141-155.
- Hall, J.A., T.C. Cantley, J.M. Galvin, B.N. Day, R.V. Anthony. 1992. Influence of ovarian steroids on relaxin-induced uterine growth in ovariectomized gilts. Endocrinol. 130:3159-3160.
- Trout, W.E., J.A. Hall, M.L. Stallings-Mann, J.M. Galvin, R.V. Anthony, R.M. Roberts. 1992. Steroid regulation of the synthesis and secretion of retinol-binding protein by the uterus of the pig. Endocrinol. 130:2557-2564.
- Caldwell, R.W., R.A. Bellows, J.A. Hall, R.V. Anthony. 1990. Administration of porcine relaxin to beef heifers four or seven days prepartum. J. Reprod. Fertil.
- Hall, J.A., T.C. Cantley, B.N. Day, R.V. Anthony. 1990. Uterotropic actions fo relaxin in prepubertal gilts. Biol. Reprod. 42:769-774.
- Weems, C.W., Y.S. Weems, J. Hall, B. DeGroot, C.N. Lee, D.L. Vincent. 1989. Effects of chronic infucion of dibutyryl-cAMP or adenosine on luteal function in sheep. Endocrinol. Japon. 36:439-447.
- Hall, J.A., E.M. Meisterling, P.E. Lewis, R.A. Dailey. 1989. Formation of ovarian cysts and copora lutea after treatment with anti-histamine or indomethacin in prepubertal gilts. Biol. Reprod. 40:565-569.

## **NON-REFEREED PUBLICATIONS:**

- Moll, F., S. Lin, J. Hall. 1994 (June). Dendrimers: An enhancement of core technology for Stratus assays. In: *Tabline*, Technical Bulletin for Stratus Immunochemistry Analyzer.



Hall, J. 1994 (August). Dendrimer Technology: Expanded flexibility of assay format capabilities for Stratus instrumentation. In: *Tabline*, Technical Bulletin for Stratus Immunochemistry Analyzer.

Caldwell, R.W., R.A. Bellows, J.A. Hall, R.V. Anthony. 1990. Administration of porcine relaxin to beef heifers four or seven days prepartum. In: 1989 Missouri Beef Cattle Report, Univ. Missouri.

Dailey, R.A., J.A. Hall, E.M. Meisterling. 1987. What causes ovarian cysts? In: Inquiry- 1987, West Virginia University Annual Research Report.

#### **SELECTED ABSTRACTS:**

Bonnar, P., M. Vasily-Cioffi, C. Chiodini, J. Bonapace, L. Baker, J. Hall. 1999. Development of a chemiluminescent assay for detection of toxoplasma IgG. Annual Meeting Amer. Soc. Micro.

Sliwinski, G., S. Bonapace, J. Green, F. Kenny, L. Baker, J. Hall. 1999. Development of a chemiluminescent assay for detection of toxoplasma IgM. Annual Meeting Amer. Soc. Micro.

Hall, J., P. Bonnar, G. Sliwinski, L. Baker. 1999. Development of toxoplasma serology assays for the ACS:Centaur. Annual Meeting Int. Fed. Clin. Chem.

Hall, J., G. Sliwinski, M. Lynch, L. Baker. 1998. An automated chemiluminescent immunoassay system for quantitative determination of specific antibodies to hepatitis B surface antigen. Annual Meeting Oak Ridge Conference.

Sliwinski, G., B. Rojik, D. Deslauriers, L. Baker, J. Hall. 1998. Development of a quantitative anti-HBs chemiluminescent assay for the ACS:Centaur immunoassay analyzer. Annual Meeting Amer. Soc. Microbiol.

Kenny, F., A. Saadat, T. Reidy, L. Baker, J. Hall. 1998. Development of an anti-HBc (IgM) chemiluminescent assay for the ACS:Centaur immunoassay analyzer. Annual Meeting Amer. Soc. Microbiol.

Hall, J., K. Broadwell, A. McKinnon, D. Masi, D. Bedard, L. Baker. 1998. Development of a hepatitis B surface antigen (HBsAg) chemiluminescent assay for the ACS:Centaur immunoassay analyzer. Annual Meeting Amer. Soc. Microbiol.

Bonnar, P., M. Cioffi, P. Wunderli, L. Baker, J. Hall. 1998. Development of a confirmatory chemiluminescent assay for hepatitis B surface antigen (HBsAg) for the ACS:Centaur. Annual Meeting Amer. Soc. Microbiol.

Wunderli, P., R. Drolet, L. Peters, T. Reidy, N. Masillo, M. Jackson, L. Baker, J. Hall. 1998. Development of an anti-HBc (total) chemiluminescent assay for the ACS:Centaur immunoassay analyzer. Annual Meeting Amer. Soc. Microbiol.

Hall, J., P. Bonnar, F. Kenny, G. Sliwinski, P. Wunderli, L. Baker. 1998. Performance of chemiluminescent immunoassays for HBV markers on the automated ACS:Centaur system. Annual Meeting Amer. Assoc. Clin. Chem.

Gammie, A., S.A. Ball, G. Sliwinski, L. Baker, C. Fang, J. Hall. 1998. Reliable quantitative measurement of anti-HBs by an automated chemiluminescent immunoassay system. Annual Meeting UK Virol. Symp.

Hall, J., G. Sliwinski, J. Green, S. Bonapace, J. Bonapace, L. Baker. 1998. HBsAg functionality affects the quantitative measurement of anti-HBs. Annual Meeting Eur. Soc. Clin. Virol.

- Wunderli, P., F. Kenny, R. Drolet, L. Peters, L. Baker, J. Hall. 1998. Importance of IgM detection in early seroconversion sensitivity of total anti-HBc assays. Annual Meeting Eur. Soc. Clin. Virol.
- Sliwinski, G., K. Vellieux, H. Kirchick, J. Hall. 1996. Development of chemiluminescent IgM anti-HBc assay for ACS instrumentation. Annual Meeting Amer. Soc. Microbiol.
- Hall, J.A., J. Cobb, F. Moll, K. Yu, J. Conde, S. Lin, P. Singh, R. Bauer. 1994. CK-MB assay performance for Stratus instrumentation using dendrimer technology. Annual Meeting Amer. Assoc. Clin. Chem.
- Hall, J.A., A.D. Okonmah, D.J. Stade, F. Boches, H.J. Kirchick. 1992. Development of IgG class capture for Stratus radial partitioning enzyme-immunoassay (RPEIA) technology. Annual Meeting Amer. Assoc. Clin. Chem.
- Hall, J.A., T.C. Cantley, J.M. Galvin, B.N. Day, R.V. Anthony. 1991. Influence of relaxin on ovarian steroid-induced uterine growth in ovariectomized gilts. Annual Meeting Endocrin. Soc.
- Hall, J.A., B.N. Day, R.V. Anthony. 1991. Influence of steroids on relaxin-induced changes in distensibility and composition of the cervix in gilts. Biol. Reprod. (Supp. 1) 44:446.
- Trout, W.E., J.A. Hall, R.V. Anthony. 1991. Uterine retinol-binding proteins of the pig: expression during pregnancy and induction by conceptus estrogen. Biol. Reprod. (Supp. 1) 44:363.
- Hall, J.A., T.C. Cantley, B.N. Day, R.V. Anthony. 1989. Uterotropic actions of relaxin in prepubertal gilts. Biol. reprod. (Supp. 1) 40:453.
- Cooper, D.A., J.A. Hall, E.K. Inskeep, R.A. dailey. 1988. Functional characteristics of induced ovarian cysts in prepubertal gilts. J. Anim. Sci. (Supp. 1) 66:518.
- Hall, J.A., E.M. Meisterling, R.A. Dailey. 1987. Effect of indomethacin on cyst formation and ovulation in prepubertal gilts. Biol. Reprod. (Supp. 1) 36:95.
- Hall, J.A., R.A. Dailey, E.K. Inskeep, P.E. Lewis. 1987. Effect of the corpus luteum of pregnancy on ovulation and luteal function in early post-partum ewes. J. Anim. Sci. (Supp. 1) 65:61.
- Hall, J.A., E.M. Meisterling, R.A. Dailey. 1986. Effect of H1 and H2 receptor antagonists on cyst formation in prepubertal gilts. J. Anim. Sci. (Supp. 1) 63:361.
- Hall, J.A. V.I. McCormick, P.E. Lewis. 1985. Effect of progestin pretreatment on gonadal steroid production and luteal activity in seasonally anestrous ewes. J. Anim. Sci. (Supp. 1) 61:135.
- Keisler, D.H., J.A. Hall, P.E. Lewis. 1983. The effect of ovariectomy on preovulatory surge of LH induced by pulsatile delivery of LH into prepubertal ewe lambs. Biol. Reprod. (Supp. 1) 28:95.



Search for product name or  
catalogue number

Advanced Search

## Search Results

| Cat No    | Product   | Pack   | Price           |
|-----------|---|--------|-----------------|
| 0100-0037 | <u>Lipoteichoic Acid (LTA)</u><br>antibody, mouse clone<br>BGN/11/40?? (IgG3): purified,<br>ELISA 1/5000 (50%B) ELISA >1/250K (EPT) | 200 µg | £155 \$269 €282 |

[About Us](#)

[Contact Us](#)

[Catalogue Info](#)

[New Products](#)

[More Info](#)

[Custom R&D Services](#)

[Distributed Products](#)

[View Your Cart](#)

[Home](#)

[Links](#)

[company profile](#) | [contact us](#) | [catalogue](#) | [request information](#)  
[view cart](#) | [help](#) | [home](#) | [terms & conditions](#)

copyright 2003 Biogenesis

→ HY-CULT BIOTECH (THE NETHERLANDS)

|                      |  |           |        |
|----------------------|--|-----------|--------|
| HM2090<br><b>NEW</b> | Mouse monoclonal antibody against Human L-Ficolin, Clone GN4 | IA, IP, W | 100 µg |
| HM2091<br><b>NEW</b> | Mouse monoclonal antibody against Human L-Ficolin, Clone GN5 | IA, IP, W | 100 µg |
| <b>*T</b>            |  |           |        |

### LPS, MICROBIAL TOXINS AND RELATED PROTEINS:

| Cat. no.            | Product   | Applications | Quantity                       |
|---------------------|---|--------------|--------------------------------|
| HM2045              | Mouse monoclonal antibody against KDO, Clone 20                     | IA, W        | >200 µg/ml<br>1 ml supernatant |
| HM2046              | Mouse monoclonal antibody against free Lipid A, Clone 43            | IA, W        | >200 µg/ml<br>1 ml supernatant |
| HM2048              | Mouse monoclonal antibody against Lipoteichoic Acid (LTA), Clone 55 | IA, W        | >200 µg/ml<br>1 ml supernatant |
| HM6001              | Mouse monoclonal antibody against LPS core, Clone WN1 222-5         | B, IA, P, W  | 100 µg                         |
| HM2047              | Mouse monoclonal antibody against Polymyxin B (PMB), Clone 45       | IA, W        | >200 µg/ml<br>1 ml supernatant |
| HC4020              | Polymyxin B (PMB)   | B, IA        | 10 <sup>5</sup> units          |
| HC4021              | Polymyxin B (PMB), biotinylated                                     | IA           | 0.2 ml                         |
| HC4022              | Polymyxin B (PMB), HRP conjugated                                   | IA           | 0.2 ml                         |
| HM6002              | Mouse monoclonal antibody against TSST-1, Clone Mab5                | B, IA, W     | 100 µg                         |
| HK506<br><b>NEW</b> | TSST-1 ELISA  | IA           | Under development              |
| <b>*T</b>           |   |              |                                |

### SAA (Serum Amyloid A)

| Cat. no.             | Product  | Applications | Quantity |
|----------------------|--|--------------|----------|
| HM2100<br><b>NEW</b> | Mouse monoclonal antibody against Human SAA<br>Clone Reu86.1 | F, IA, P     | 100 µg   |
| HM2101<br><b>NEW</b> | Mouse monoclonal antibody against Human SAA<br>Clone Reu86.5 | F, IA, P     | 100 µg   |
| <b>*TOP</b>          |  |              |          |


### SLPI (Secretory Leucocyte Protease Inhibitor):

Secretory leucocyte protease inhibitor (SLPI; also known as antileukoprotease or ALP) is a 11.7 kDa cati inhibitor of neutrophil elastase and to a lesser extent of cathepsin G. It is produced by epithelial cells in tl lung, skin and other organs and by PMN and (in mice) by macrophages.

In addition to its protease inhibitory properties that may serve to protect against proteolytic injury, it was

Figure 5

## Gram-Positive Bacterial Recognition



## INFECTIOUS DISEASE ANTIBODIES

**E. coli K99 pili (LP)**

#18401-18404

(4 clones)

**Price:** \$235/mg or  
\$235/ml**Host:** Mouse

These monoclonal antibodies recognize *E. coli* K99 pili and do not cross-react with K99-negative *E. coli*, *E. coli* K88ac, *E. coli* 987P, *Proteus mirabilis*, *Proteus vulgaris*, *Pseudomonas aeruginosa*, *Enterobacter aerogenes*, *Citrobacter freundii*, or *Klebsiella pneumoniae*. These antibodies may be used in immunoassays to identify K99-positive *E. coli*. Available as ascites, purified antibody, and in a Library Pack.

> **Endotoxin (LP)**

#15301-15309

(9 clones)

**Price:** \$235/mg or  
\$235/ml**Host:** Mouse

These monoclonal antibodies recognize endotoxin produced by gram-negative bacteria. Clones differ in degrees of cross-reactivity with a panel of gram-negative endotoxins including *E. coli*, *Klebsiella pneumoniae*, *Shigella sonnei*, *Salmonella typhimurium*, *Enterobacter aerogenes*, *Serratia marcescens*, *Proteus mirabilis*, *Proteus vulgaris*, *Acinetobacter calcoaceticus*, and *Pseudomonas aeruginosa*. Available as ascites, purified antibody, and in a Library Pack.

**Entamoeba histolytica (LP)**

#15000, 15001, 15010, 15020, 15030

(5 clones)

**Price:** \$235/mg or  
\$235/ml**Host:** Mouse

These monoclonal antibodies recognize *Entamoeba histolytica* trophozoites and cysts. These antibodies may be used in immunoassays to detect *E. histolytica*. Available as ascites, purified antibody, and in a Library Pack.

**Eotaxin**

#1123

**Price:** \$210/100 ug**Host:** Rabbit

This antibody recognizes human eotaxin, an eosinophil chemoattractant. The eotaxin receptor CCR3 is required for HIV-1 entry into target cells, and eotaxin inhibits infection by HIV-1. A monomer band of 9 kD and a homodimer band of 18 kD are detected in immunoblots. The epitope for this antibody is in amino acids 83-97 of human eotaxin.

# MONOTOPE™

(Mouse)  
Monoclonal Antibodies\*  
Continued from previous page

| AGENT                         | SPECIFICITY  | PURIFIED<br>(100 UG) | FITC CONJ.<br>(100 UG) |
|-------------------------------|--------------|----------------------|------------------------|
| Dengue virus                  | group        | 5161 \$130           |                        |
| Diphtheria toxin              | —            | 7801 \$130           |                        |
| Diphtheria toxin              | A subunit    | 7811 \$130           |                        |
| E. coli                       | common       | 1011 \$130           |                        |
| E. coli                       | O157         | 1031 \$145           | 1033 \$170             |
| E. coli                       | O157         | 1041 \$145           |                        |
| E. coli                       | O157         | 1051 \$160           |                        |
| E. coli                       | K99          | 6911 \$160           |                        |
| > Endotoxin                   | gram neg.    | 6901 \$160           |                        |
| Enterobacteriaceae            | common       | 1121 \$130           |                        |
| Epstein-Barr virus            | EBNA-1       | 0211 \$130           |                        |
| Epstein-Barr virus            | EBNA-1       | 0216 \$130           |                        |
| Epstein-Barr virus            | gp 250/350   | 0221 \$130           | 0223 \$165             |
| Epstein-Barr virus            | VCA          | 0231 \$130           |                        |
| Epstein-Barr virus            | VCA-gp       | 0241 \$130           |                        |
| Epstein-Barr virus            | EA-R         | 0251 \$130           |                        |
| Epstein-Barr virus            | EA-D         | 0261 \$130           |                        |
| Feline Calicivirus            | —            | 7401 \$130           |                        |
| Feline Calicivirus            | —            | 7402 \$130           |                        |
| Feline Immunodeficiency virus | FIV p24      | 7301 \$130           |                        |
| Feline Immunodeficiency virus | FIV p24      | 7304 \$130           |                        |
| Feline Immunodeficiency virus | protease     | 7305 \$130           |                        |
| Feline Immunodeficiency virus | FIV p24      | 7306 \$130           |                        |
| Feline Immunodeficiency virus | FIV p24      | 7307 \$130           |                        |
| Feline Immunodeficiency virus | FIV envelope | 7312 \$130           |                        |
| Feline Inf. Peritonitis virus | Coronavirus  | 7321 \$130           |                        |
| Feline Leukemia virus         | FeLv p27     | 7201 \$130           |                        |
| Feline Leukemia virus         | FeLv gp7O    | 7202 \$130           |                        |
| Feline Leukemia virus         | FeLv gp7O    | 7203 \$130           |                        |
| Feline Leukemia virus         | FeLv gp7O    | 7204 \$130           |                        |
| Feline Leukemia virus         | FeLv gp7O    | 7205 \$130           |                        |
| Feline Leukemia virus         | FeLv gp7O    | 7206 \$130           |                        |
| Feline Leukemia virus         | FeLv gp7O    | 7207 \$130           |                        |
| Feline Leukemia virus         | FeLv gp7O    | 7208 \$130           |                        |
| Feline Leukemia virus         | FeLv gp7O    | 7209 \$130           |                        |
| Feline Leukemia virus         | FeLv p15e    | 7210 \$130           |                        |
| Feline Leukemia virus         | FeLv p15e    | 7211 \$130           |                        |
| Feline Leukemia virus         | FeLv p15e    | 7212 \$130           |                        |
| Feline Leukemia virus         | FeLv p15e    | 7213 \$130           |                        |
| Feline Leukemia virus         | FeLv p15e    | 7214 \$130           |                        |
| Feline Leukemia virus         | FeLv p15e    | 7215 \$130           |                        |
| Feline Rhinotracheitis virus  | —            | 7531 \$130           |                        |
| Giardia lamblia               | trophozoites | 6211 \$130           |                        |
| Helicobacter pylori           | —            | 6611 \$130           |                        |

E. coli

Epstein-Barr  
virus

.....ViroStat • 207-856-6620 Fax 207-856-6864

\* For Research Use Only — Not for Use in Diagnostic Procedures.

↳ HY-CULT BIOTECH (THE NETHERLANDS)

|                      |  |           |        |
|----------------------|--|-----------|--------|
| HM2090<br><b>NEW</b> | Mouse monoclonal antibody against Human L-Ficolin, Clone GN4 | IA, IP, W | 100 µg |
| HM2091<br><b>NEW</b> | Mouse monoclonal antibody against Human L-Ficolin, Clone GN5 | IA, IP, W | 100 µg |
| <b>*T</b>            |  |           |        |

### LPS, MICROBIAL TOXINS AND RELATED PROTEINS:

| Cat. no.            | Product   | Applications | Quantity                       |
|---------------------|---|--------------|--------------------------------|
| HM2045              | Mouse monoclonal antibody against KDO, Clone 20                     | IA, W        | >200 µg/ml<br>1 ml supernatant |
| HM2046              | Mouse monoclonal antibody against free Lipid A, Clone 43            | IA, W        | >200 µg/ml<br>1 ml supernatant |
| HM2048              | Mouse monoclonal antibody against Lipoteichoic Acid (LTA), Clone 55 | IA, W        | >200 µg/ml<br>1 ml supernatant |
| HM6001              | Mouse monoclonal antibody against LPS core, Clone WN1 222-5         | B, IA, P, W  | 100 µg                         |
| HM2047              | Mouse monoclonal antibody against Polymyxin B (PMB), Clone 45       | IA, W        | >200 µg/ml<br>1 ml supernatant |
| HC4020              | Polymyxin B (PMB)   | B, IA        | 10 <sup>5</sup> units          |
| HC4021              | Polymyxin B (PMB), biotinylated                                     | IA           | 0.2 ml                         |
| HC4022              | Polymyxin B (PMB), HRP conjugated                                   | IA           | 0.2 ml                         |
| HM6002              | Mouse monoclonal antibody against TSST-1, Clone Mab5                | B, IA, W     | 100 µg                         |
| HK506<br><b>NEW</b> | TSST-1 ELISA  | IA           | Under development              |
| <b>*T</b>           |   |              |                                |

### SAA (Serum Amyloid A)

| Cat. no.             | Product  | Applications | Quantity |
|----------------------|--|--------------|----------|
| HM2100<br><b>NEW</b> | Mouse monoclonal antibody against Human SAA<br>Clone Reu86.1 | F, IA, P     | 100 µg   |
| HM2101<br><b>NEW</b> | Mouse monoclonal antibody against Human SAA<br>Clone Reu86.5 | F, IA, P     | 100 µg   |
| <b>*TOP</b>          |  |              |          |

### SLPI (Secretory Leucocyte Protease Inhibitor):

Secretory leucocyte protease inhibitor (SLPI; also known as antileukoprotease or ALP) is a 11.7 kDa cati inhibitor of neutrophil elastase and to a lesser extent of cathepsin G. It is produced by epithelial cells in tl lung, skin and other organs and by PMN and (in mice) by macrophages.

In addition to its protease inhibitory properties that may serve to protect against proteolytic injury, it was



Figure 6

## Gram-Negative Bacterial Recognition

